arXiv:1112.3429v2 [math.RT] 6 Aug 2012

NON-ORTHOGONAL GEOMETRIC REALIZATIONS OF COXETER GROUPS

FU, XIANG

School of Mathematics and Statistics
University of Sydney, NSW 2006, Australia

`xifu9119@mail.usyd.edu.au`
`xiangf@maths.usyd.edu.au`

Preliminary version, August 7, 2012

Abstract. We define in an axiomatic fashion a *Coxeter datum* for an arbitrary Coxeter group W. Such Coxeter datum will specify a pair of reflection representations of W which are not assumed to be embeddings of W into the orthogonal group of any real vector space. These non-orthogonal representations then give rise to a pair of inter-related root systems that generalize the usual orthogonal root systems of Coxeter groups. We obtain comparison results between these non-orthogonal root systems and the standard (orthogonal) root systems. Further, we study cones analogous to the Tits cones.

1. Introduction

For an arbitrary Coxeter system (W, R) in the sense of [13] or [17], it is well known that W can be embedded into the orthogonal group of a certain bilinear form $(\, , \,)$ on a vector space V over the real field $\mathbb{R}$. The root system of W in V is a certain W-stable subset of V whose elements correspond to the reflections in W. If the bilinear form $(\, , \,)$ is non-degenerate then (of course) V is W-isomorphic to its algebraic dual $\mathrm{Hom}(V, \mathbb{R})$, but since the form is not always non-degenerate it is sometimes useful to study both the representation of W on V and the contragredient representation of W on $\mathrm{Hom}(V, \mathbb{R})$. This motivates the approach taken in this paper, in which we consider a pair of real vector spaces V_1 and V_2 linked by a W-equivariant bilinear pairing satisfying a few extra conditions (which are guaranteed to hold in the case that $V_1 = V$ and $V_2 = \mathrm{Hom}(V, \mathbb{R})$). We show that in this situation W embeds (faithfully) in the general linear groups of each of V_1 and V_2, with both images being generated by involutions. The classical theory of (orthogonal) geometric realizations can be recovered from this construction as a special case. We define and study generalized

---

2010 *Mathematics Subject Classification.* 20F55 (20F10, 20F65).

Key words and phrases. Coxeter groups, root systems, Tits cone, dominance.

root systems arising from such non-orthogonal geometric realizations of Coxeter systems, and compare them with the root systems arising from the standard geometric realizations. It turns out that it is natural to consider root systems in both V_1 and V_2 simultaneously; these are in W-equivariant bijective correspondence with each other, and are also closely related to the classical root system of W in the sense of [2] and [17].

2. Paired reflection representations

Let S be an arbitrary set in which each unordered pair $\{s, t\}$ of elements is assigned an $m_{st} \in \mathbb{Z} \cup \{\infty\}$, subject to the conditions that $m_{ss} = 1$ (for all s in S), and $m_{st} \geq 2$ (for all distinct s, t in S). Suppose that V_1 and V_2 are vector spaces over the real field $\mathbb{R}$, and suppose that there exists a bilinear map $\langle\ ,\ \rangle : V_1 \times V_2 \to \mathbb{R}$ and sets $\Pi_1 = \{\ \alpha_s \mid s \in S\ \} \subseteq V_1$ and $\Pi_2 = \{\ \beta_s \mid s \in S\ \} \subseteq V_2$ such that the following conditions hold:

(C1) $\langle \alpha_s, \beta_s \rangle = 1$, for all $s \in S$;
(C2) $\langle \alpha_s, \beta_t \rangle \leq 0$, for all distinct $s, t \in S$;
(C3) for all s, $t \in S$,

$$\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle = \begin{cases} \cos^2(\pi/m_{st}) & \text{if } m_{st} \neq \infty, \\ \gamma^2, \text{ for some } \gamma \geq 1 & \text{if } m_{st} = \infty; \end{cases}$$

(C4) $\langle \alpha_s, \beta_t \rangle = 0$ if and only if $\langle \alpha_t, \beta_s \rangle = 0$, for all $s, t \in S$;
(C5) $\sum_{s \in S} \lambda_s \alpha_s = 0$ with $\lambda_s \geq 0$ for all s implies $\lambda_s = 0$ for all s, and $\sum_{s \in S} \lambda_s \beta_s = 0$ with $\lambda_s \geq 0$ for all s implies $\lambda_s = 0$ for all s.

Note that (C3) and (C4) together imply that $\langle \alpha_s, \beta_t \rangle$ and $\langle \alpha_t, \beta_s \rangle$ are zero if and only if $m_{st} = 2$. We can also express (C5) more compactly as $0 \notin \mathrm{PLC}(\Pi_1)$ and $0 \notin \mathrm{PLC}(\Pi_2)$, where $\mathrm{PLC}(A)$ (the *positive linear combinations of* A) of any set A is defined to be

$$\{ \sum_{a \in A} \lambda_a a \mid \lambda_a \geq 0 \text{ for all } a \in A, \text{ and } \lambda_{a'} > 0 \text{ for some } a' \in A \}.$$

Definition 2.1. In the above situation, if conditions (C1) to (C5) are satisfied then we call $\mathscr{C} = (S, V_1, V_2, \Pi_1, \Pi_2, \langle\ ,\ \rangle)$ a *Coxeter datum.* The m_{st} $(s, t \in S)$ are called the *Coxeter parameters* of $\mathscr{C}$.

Following [6], we call $\mathscr{C}$ *free* if both Π_1 and Π_2 are linearly independent. Throughout this paper, $\mathscr{C} = (S, V_1, V_2, \Pi_1, \Pi_2, \langle\ ,\ \rangle)$ will be a fixed Coxeter datum with Coxeter parameters m_{st}, unless otherwise stated. We stress that, in general, $\mathscr{C}$ is not required to be free.

Remark 2.2. In a Coxeter datum $\mathscr{C}$, if we take $V_2 = \mathrm{Hom}(V_1, \mathbb{R})$ and take $\langle\, ,\, \rangle$ to be the natural pairing on $V_1 \times \mathrm{Hom}(V_1, \mathbb{R})$, then the conditions (RB1), (RB2) and (RB3) required in the definition of the *root data* of [6] are automatically satisfied in $\mathscr{C}$ (though, here we have scaled each element of Π_1 and Π_2 by a factor of $\frac{1}{\sqrt{2}}$). Thus the root data in the sense of [6] are special cases of Coxeter datum defined here. Further, if we assume that $\mathscr{C}$ is free then we recover a set of root data as given in [12], whereas if we assume that $\langle \alpha_s, \beta_t \rangle \in \mathbb{Z}$ for all $s, t \in S$ then we recover a set of root data as given in [21].

Definition 2.3. Given a Coxeter datum $\mathscr{C} = (S, V_1, V_2, \Pi_1, \Pi_2, \langle\ ,\ \rangle)$, for each $s \in S$, let $\rho_{V_1}(s)$ and $\rho_{V_2}(s)$ be the linear transformations on V_1 and V_2 defined by

$$\rho_{V_1}(s)(x) = x - 2\langle x, \beta_s \rangle \alpha_s$$

for all $x \in V_1$, and

$$\rho_{V_2}(s)(y) = y - 2\langle \alpha_s, y \rangle \beta_s$$

for all $y \in V_2$. For each $i \in \{1, 2\}$ let $R_i(\mathscr{C}) = \{\, \rho_{V_i}(s) \mid s \in S \,\}$, and let $W_i(\mathscr{C})$ be the subgroup of $\mathrm{GL}(V_i)$ generated by R_i.

For each $s \in S$, it is readily checked that $\rho_{V_1}(s)$ and $\rho_{V_2}(s)$ are involutions with $\rho_{V_1}(s)(\alpha_s) = -\alpha_s$ and $\rho_{V_2}(s)(\beta_s) = -\beta_s$. Further, we have:

Proposition 2.4. *Let $x \in V_1$ and $y \in V_2$. Then for all $s \in S$,*

$$\langle \rho_{V_1}(s)(x), \rho_{V_2}(s)(y) \rangle = \langle x, y \rangle.$$

□

Though $\mathscr{C}$ lacks freeness in general, nevertheless, conditions (C1), (C2) and (C5) of the definition of a Coxeter datum together yield that:

Lemma 2.5. *For each $s \in S$ we have $\alpha_s \notin \mathrm{PLC}(\Pi_1 \setminus \{\alpha_s\})$, and $\beta_s \notin \mathrm{PLC}(\Pi_2 \setminus \{\beta_s\})$. In particular, for distinct $s, t \in S$, the set $\{\alpha_s, \alpha_t\}$ is linearly independent, and so is $\{\beta_s, \beta_t\}$.* □

For each $i \in \{1, 2\}$, the above lemma yields that $\rho_{V_i}(s) \neq \rho_{V_i}(t)$ whenever $s, t \in S$ are distinct. The following lemma, readily obtained from direct calculations, summarizes a few useful results:

Lemma 2.6. *(i) Suppose that $s, t \in S$ such that $m_{st} \notin \{1, \infty\}$, and let $\theta = \pi / m_{st}$. If $m_{st} \neq 2$ then for each $n \in \mathbb{N}$,*

$$(\rho_{V_1}(s)\rho_{V_1}(t))^n(\alpha_s) = \frac{\sin(2n+1)\theta}{\sin\theta}\alpha_s + \frac{-\cos\theta}{\langle \alpha_t, \beta_s \rangle}\frac{\sin(2n\theta)}{\sin\theta}\alpha_t$$

and

$$(\rho_{V_2}(s)\rho_{V_2}(t))^n(\beta_s) = \frac{\sin(2n+1)\theta}{\sin\theta}\beta_s + \frac{-\cos\theta}{\langle \alpha_s, \beta_t \rangle}\frac{\sin(2n\theta)}{\sin\theta}\beta_t.$$

While if $m_{st} = 2$ then for each $n \in \mathbb{N}$,

$$(\rho_{V_1}(s)\rho_{V_1}(t))^n(\alpha_s) = (-1)^n\alpha_s, \quad \textit{and} \quad (\rho_{V_2}(s)\rho_{V_2}(t))^n(\beta_s) = (-1)^n\beta_s.$$

(ii) Suppose that $s, t \in S$ such that $m_{st} = \infty$. Define $\theta = \cosh^{-1}(\gamma)$, where $\gamma = \sqrt{\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle}$. Then for each $n \in \mathbb{N}$,

$$(\rho_{V_1}(s)\rho_{V_1}(t))^n(\alpha_s) = \begin{cases} \frac{\sinh(2n+1)\theta}{\sinh\theta}\alpha_s + \frac{-\gamma}{\langle \alpha_t, \beta_s \rangle}\frac{\sinh(2n\theta)}{\sinh\theta}\alpha_t, & \text{if } \theta \neq 0, \\ (2n+1)\alpha_s - \frac{2n\gamma}{\langle \alpha_t, \beta_s \rangle}\alpha_t, & \text{if } \theta = 0; \end{cases}$$

and

$$(\rho_{V_2}(s)\rho_{V_2}(t))^n(\beta_s) = \begin{cases} \frac{\sinh(2n+1)\theta}{\sinh\theta}\beta_s + \frac{-\gamma}{\langle \alpha_s, \beta_t \rangle}\frac{\sinh(2n\theta)}{\sinh\theta}\beta_t, & \text{if } \theta \neq 0, \\ (2n+1)\beta_s - \frac{2n\gamma}{\langle \alpha_s, \beta_t \rangle}\beta_t, & \text{if } \theta = 0. \end{cases}$$

□

Remarks 2.7. Let $\mathrm{ord}(\rho_{V_i}(s)\rho_{V_i}(t))$ denote the order of $\rho_{V_i}(s)\rho_{V_i}(t)$ in $\mathrm{GL}(V_i)$, for each $i \in \{1, 2\}$. Then:
(i) Lemma 2.6 (i) yields that $\mathrm{ord}(\rho_{V_i}(s)\rho_{V_i}(t)) \geq m_{st}$ when $m_{st} \neq \infty$. Indeed, in the subspace with basis $\{\alpha_s, \alpha_t\}$ the following m_{st} elements

$$\alpha_s, (\rho_{V_1}(s)\rho_{V_1}(t))\alpha_s, (\rho_{V_1}(s)\rho_{V_1}(t))^2\alpha_s, \ldots, (\rho_{V_1}(s)\rho_{V_1}(t))^{m_{st}-1}\alpha_s$$

are all distinct, and the same holds in the subspace with basis $\{\beta_s, \beta_t\}$.
(ii) In a similar way as the above, it follows from Lemma 2.6 (ii) that $\mathrm{ord}(\rho_{V_i}(s)\rho_{V_i}(t)) = \infty$ when $m_{st} = \infty$.

The above observations naturally lead to the following:

Proposition 2.8. *Suppose that $s, t \in S$. Then for each $i \in \{1, 2\}$,*

$\rho_{V_i}(s)\rho_{V_i}(t)$ has order m_{st} in $\mathrm{GL}(V_i)$.

Proof. We give a proof that $\rho_{V_1}(s)\rho_{V_1}(t)$ has order m_{st} in $\mathrm{GL}(V_1)$ below, and we stress that the same argument will hold for $\rho_{V_2}(s)\rho_{V_2}(t)$ in $\mathrm{GL}(V_2)$. Observe that we only need to consider the cases when $m_{st} \notin \{1, \infty\}$, for the statement of the proposition follows readily from Remark 2.7 (ii) and the fact that each $\rho_{V_1}(s)$, for all $s \in S$, is an involution. Next let $\alpha \in V_1$ be arbitrary. Then

$$\begin{aligned} (\rho_{V_1}(s)\rho_{V_1}(t))(\alpha) &= \rho_{V_1}(s)(\alpha - 2\langle \alpha, \beta_t \rangle \alpha_t) \\ &= \alpha - 2\langle \alpha, \beta_s \rangle \alpha_s - 2\langle \alpha, \beta_t \rangle (\alpha_t - 2\langle \alpha_t, \beta_s \rangle \alpha_s) \\ &= \alpha + (4\langle \alpha, \beta_t \rangle \langle \alpha_t, \beta_s \rangle - 2\langle \alpha, \beta_s \rangle)\alpha_s - 2\langle \alpha, \beta_t \rangle \alpha_t. \end{aligned} \tag{2.1}$$

If $m_{st} = 2$, then (2.1) yields that

$$(\rho_{V_1}(s)\rho_{V_1}(t))(\alpha) = (\rho_{V_1}(t)\rho_{V_1}(s))(\alpha) = \alpha - 2\langle \alpha, \beta_s \rangle \alpha_s - 2\langle \alpha, \beta_t \rangle \alpha_t,$$

so that $\rho_{V_1}(s)$ and $\rho_{V_1}(t)$ commute. Hence $\mathrm{ord}(\rho_{V_1}(s)\rho_{V_1}(t)) = 2$ when $m_{st} = 2$, and it remains to check the case when $m_{st} > 2$. Observe that if $\alpha = \alpha_s$ and $\alpha = \alpha_t$, then (2.1) yields that

$$(\rho_{V_1}(s)\rho_{V_1}(t))(\alpha_s) = (4\cos^2(\pi/m_{st}) - 1)\alpha_s - 2\langle \alpha_s, \beta_t \rangle \alpha_t$$

and

$$(\rho_{V_1}(s)\rho_{V_1}(t))(\alpha_t) = 2\langle \alpha_t, \beta_s\rangle \alpha_s - \alpha_t.$$

Therefore the action of $\rho_{V_1}(s)\rho_{V_1}(t)$ on $\mathbb{R}\{\alpha_s, \alpha_t, \alpha\}$ may be represented by the following matrix M:

$$M = \begin{pmatrix} 4\cos^2(\pi/m_{st}) - 1 & 2\langle \alpha_t, \beta_s\rangle & 4\langle \alpha, \beta_t\rangle\langle \alpha_t, \beta_s\rangle - 2\langle \alpha, \beta_s\rangle \\ -2\langle \alpha_s, \beta_t\rangle & -1 & -2\langle \alpha, \beta_t\rangle \\ 0 & 0 & 1 \end{pmatrix}.$$

It is readily checked that M has distinct eigenvalues $e^{i\frac{2\pi}{m_{st}}}$, $e^{-i\frac{2\pi}{m_{st}}}$ and 1. Hence M has order m_{st}, and so $(\rho_{V_1}(s)\rho_{V_1}(t))^{m_{st}} = 1$ in $\mathrm{GL}(V_1)$. Finally, in view of Remark 2.7 (i), it follows that $\mathrm{ord}(\rho_{V_1}(s)\rho_{V_1}(t))$ is precisely m_{st}. $\square$

Remark 2.9. Given a Coxeter datum $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle,\rangle\,)$ with Coxeter parameters m_{st} (where $s, t \in S$), let (W, R) be a Coxeter system in the sense of [13] or [17] with $R = \{\, r_s \mid s \in S\,\}$ being a set of involutions generating W subject only to the condition that the order of the product $r_s r_t$ is m_{st} whenever s, t are in S with $m_{st} \neq \infty$. Then Proposition 2.8 yields that there are group homomorphisms $f_1 \colon W \to W_1(\mathscr{C})$ and $f_2 \colon W \to W_2(\mathscr{C})$ satisfying $f_1(r_s) = \rho_{V_1}(s)$ and $f_2(r_s) = \rho_{V_2}(s)$ for all $s \in S$.

The principal result of this section is the following:

Theorem 2.10. *Let (W, R), f_1 and f_2 be as the above. Then f_1 and f_2 are isomorphisms; that is, $(W_1(\mathscr{C}), R_1(\mathscr{C}))$ and $(W_2(\mathscr{C}), R_2(\mathscr{C}))$ are both Coxeter systems isomorphic to (W, R).*

For each $i \in \{1, 2\}$, since $W_i(\mathscr{C})$ is generated by the elements of $R_i(\mathscr{C})$, it follows readily that each f_i is surjective. Thus only the injectivity of f_i needs to be checked. Before we can do so, a few elementary results are needed. First we have a result easily obtained from the formulas in Lemma 2.6:

Lemma 2.11. *Suppose that $s, t \in S$, and let n be an integer such that $0 \le n < m_{st}$. Write*

$$\underbrace{\cdots \rho_{V_1}(t)\rho_{V_1}(s)\rho_{V_1}(t)}_{n \text{ factors}}(\alpha_s) = \lambda_n \alpha_s + \mu_n \alpha_t$$

and

$$\underbrace{\cdots \rho_{V_1}(s)\rho_{V_1}(t)\rho_{V_1}(s)}_{n \text{ factors}}(\alpha_t) = \lambda'_n \alpha_s + \mu'_n \alpha_t.$$

Then all four constants λ_n, μ_n, λ'_n and μ'_n are non-negative. $\square$

Remark 2.12. The same argument applies equally well if we replace in the above lemma, respectively, $\rho_{V_1}(s)$, $\rho_{V_1}(t)$, α_s and α_t by $\rho_{V_2}(s)$, $\rho_{V_2}(t)$, β_s and β_t.

Let W, f_1 and f_2 be as in Remark 2.9. Then f_1 and f_2 give rise to W-actions on V_1 and V_2 in the following way: $wx = (f_1(w))(x)$ for all $w \in W$ and $x \in V_1$, and $wy = (f_2(w))(y)$ for all $w \in W$ and $y \in V_2$. Let $\ell \colon W \to \mathbb{N}$ be the *length* function of W with respect to R. For $w \in W$ we say that an expression of the form $w = r_{s_1} \cdots r_{s_l}$ (where $s_1, \dots s_l \in S$) *reduced* if $\ell(w) = l$. For any $w \in W$, an easy induction on $\ell(w)$ yields the following extension to Proposition 2.4:

Lemma 2.13. *Given a Coxeter datum* $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle , \rangle\,)$, *and let* W *be as in in Remark 2.9. Then* $\langle x, y\rangle = \langle wx, wy\rangle$ *for all* $w \in W$, $x \in V_1$, *and* $y \in V_2$. ☐

Using the same argument as in [15], we may deduce:

Proposition 2.14. ([15, Theorem, Lecture 1]) *Let* W *be as the above, and let* $w \in W$ *and* $s \in S$. *If* $\ell(wr_s) \geq \ell(w)$ *then* $w\alpha_s \in \mathrm{PLC}(\Pi_1)$. ☐

Now we are ready to complete the proof of Theorem 2.10:

Proof of Theorem 2.10. Suppose, for a contradiction, that the kernel of f_1 is nontrivial, and choose w in the kernel of f_1 with $w \neq 1$. Then $\ell(w) > 0$, and we may write $w = w'r_s$ for some $s \in S$ and $w' \in W$ with $\ell(w') = \ell(w) - 1$. Since $\ell(w'r_s) > \ell(w')$, Proposition 2.14 yields $w'\alpha_s \in \mathrm{PLC}(\Pi_1)$. But then

$$\alpha_s = w\alpha_s = (w'r_s)\alpha_s = w'(r_s\alpha_s) = w'(-\alpha_s) = -w'\alpha_s,$$

and hence $0 = \alpha_s + w'\alpha_s \in \mathrm{PLC}(\Pi_1)$, contradicting condition (C5) of a Coxeter datum. In an entirely similar way it can be shown that the kernel of f_2 is trivial. ☐

Let W and R be as in Remark 2.9. We call W the *abstract Coxeter group* determined by the Coxeter parameters of the Coxeter datum $\mathscr{C}$, and we call (W, R) the *abstract Coxeter system* associated with $\mathscr{C}$. Observe that Theorem 2.10 yields that the W-actions on V_1 and V_2 induced by the isomorphisms f_1 and f_2 are faithful.

3. Root Systems and Canonical Coefficients

Definition 3.1. Suppose that $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\,,\,\rangle\,)$ is a Coxeter datum, and suppose that (W, R) is the associated abstract Coxeter system.
(i) Define $\Phi_1(\mathscr{C}) = W\Pi_1 = \{\, w\alpha_s \mid w \in W, s \in S\,\}$, and similarly define $\Phi_2(\mathscr{C}) = W\Pi_2 = \{\, w\beta_s \mid w \in W, s \in S\,\}$. For each $i \in \{1, 2\}$, we call $\Phi_i(\mathscr{C})$ the *root system* of W in V_i, and its elements the *roots* of W in V_i. We call Π_i the set of *simple roots* in $\Phi_i(\mathscr{C})$, and we say that Π_i forms a *root basis* for $\Phi_i(\mathscr{C})$.
(ii) Set $\Phi_i^+(\mathscr{C}) = \Phi_i(\mathscr{C}) \cap \mathrm{PLC}(\Pi_i)$, and $\Phi_i^-(\mathscr{C}) = -\Phi_i^+(\mathscr{C})$ for each $i \in \{1, 2\}$. We call $\Phi_i^+(\mathscr{C})$ the set of *positive roots* in $\Phi_i(\mathscr{C})$, and $\Phi_i^-(\mathscr{C})$ the set of *negative roots* in $\Phi_i(\mathscr{C})$.

Given the above notations, for each $i \in \{1, 2\}$, we adopt the traditional diagrammatic description of simple roots Π_i: draw a graph that has one vertex for each $s \in S$, and join the vertices corresponding to $s, t \in S$ by an edge labelled by m_{st} if $m_{st} > 2$. The label m_{st} is often omitted if $m_{st} = 3$. Thus the diagram  corresponds to $\Pi_1 = \{\, \alpha_r, \alpha_s, \alpha_t \,\}$ and $\Pi_2 = \{\, \beta_r, \beta_s, \beta_t \,\}$. Suppose that

$$\begin{aligned} \langle \alpha_r, \beta_s \rangle &= -1/4, & \langle \alpha_s, \beta_t \rangle &= -1/6, & \langle \alpha_t, \beta_r \rangle &= -1/10, \\ \langle \alpha_s, \beta_r \rangle &= -1, & \langle \alpha_t, \beta_s \rangle &= -3/2, & \langle \alpha_r, \beta_t \rangle &= -5/2. \end{aligned}$$

Then

$$\begin{aligned} r_s\alpha_r &= \alpha_r - 2\langle \alpha_r, \beta_s \rangle \alpha_s = \alpha_r + \tfrac{1}{2}\alpha_s; \\ (r_r r_s)\alpha_r &= r_r(\alpha_r + \tfrac{1}{2}\alpha_s) = \tfrac{1}{2}\alpha_s; \\ (r_t r_r r_s)\alpha_r &= r_t(\tfrac{1}{2}\alpha_s) = \tfrac{1}{2}\alpha_s + \tfrac{1}{6}\alpha_t; \\ (r_s r_t r_r r_s)\alpha_r &= r_s(\tfrac{1}{2}\alpha_s + \tfrac{1}{6}\alpha_t) = \tfrac{1}{6}\alpha_t; \\ (r_r r_s r_t r_r r_s)\alpha_r &= r_r(\tfrac{1}{6}\alpha_t) = \tfrac{1}{6}\alpha_t + \tfrac{1}{30}\alpha_r; \\ (r_t r_r r_s r_t r_r r_s)\alpha_r &= r_t(\tfrac{1}{6}\alpha_t + \tfrac{1}{30}\alpha_r) = \tfrac{1}{30}\alpha_r. \end{aligned}$$

In particular, we notice from the above that it is possible for a nontrivial positive scalar multiple of a root to also be a root, lying in the same W-orbit as the root itself. Clearly if $w\alpha = \lambda\alpha$ where $\alpha \in \Phi_1(\mathscr{C})$ then $w^n\alpha = \lambda^n\alpha$ for all $n \in \mathbb{N}$. Since it is quite possible that $\lambda \neq \pm 1$, it follows that there could well be infinitely many non-trivial scalar multiples of α in $\Phi_1(\mathscr{C})$. Further, all roots in the W-orbit of α will possess this same property. Of course, the same situation could arise in $\Phi_2(\mathscr{C})$ as well. This is one of the features setting $\Phi_1(\mathscr{C})$ and $\Phi_2(\mathscr{C})$ apart from the classical root systems studied in [2], [13] or [17].

Similar to [17], Proposition 2.14 yields the following

Lemma 3.2. *Suppose that* $\mathscr{C} = (\, S, V_1, V_2, \Pi_1, \Pi_2, \langle\, ,\, \rangle \,)$ *is a Coxeter datum, and suppose that* (W, R) *is the associated abstract Coxeter system.* (i) $\Phi_i(\mathscr{C}) = \Phi_i^+(\mathscr{C}) \uplus \Phi_i^-(\mathscr{C})$, *for each* $i \in \{1, 2\}$, *where* $\uplus$ *denotes disjoint union.*
(ii) *If* $w \in W$ *and* $s \in S$, *then*

$$\ell(wr_s) = \begin{cases} \ell(w) + 1 & \textit{if } w\alpha_s \in \Phi_1^+(\mathscr{C}), \textit{ and } w\beta_s \in \Phi_2^+(\mathscr{C}), \\ \ell(w) - 1 & \textit{if } w\alpha_s \in \Phi_1^-(\mathscr{C}), \textit{ and } w\beta_s \in \Phi_2^-(\mathscr{C}). \end{cases}$$

□

Remark 3.3. Part (ii) of the above lemma implies that for $w \in W$ and $s \in S$, $w\alpha_s \in \Phi_1^+(\mathscr{C})$ if and only if $w\beta_s \in \Phi_2^+(\mathscr{C})$, and $w\alpha_s \in \Phi_1^-(\mathscr{C})$ if and only if $w\beta_s \in \Phi_2^-(\mathscr{C})$.

Remark 3.4. Since $\mathscr{C}$ is not assumed to be free, although we know from Lemma 3.2 (i) that each root in $\Phi_i(\mathscr{C})$ (for each $i \in \{1, 2\}$) is expressible as a linear combination of simple roots from Π_i with coefficients all being of the same sign, that expression needs not be unique. Thus the concept of *the coefficient of a simple root in a given root* is potentially ambiguous. To obtain a canonical way of expressing roots in terms of simple roots, we employ a construction similar as those of [11], [16] and [21]. We define a free Coxeter datum $\mathscr{C}'$ on the same set of Coxeter parameters as those of $\mathscr{C}$. Then both $\mathscr{C}$ and the free Coxeter datum $\mathscr{C}'$ correspond to the same abstract Coxeter system (W, R). It turns out that for each $i \in \{1, 2\}$, there exists a canonical W-equivariant bijection $\pi_i \colon \Phi_i(\mathscr{C}) \leftrightarrow \Phi_i(\mathscr{C}')$ which maps simple roots to simple roots. Since each $\Phi_i(\mathscr{C}')$ is free, there is no ambiguity of the coefficient of a simple root in any given root of $\Phi(\mathscr{C}')$, and this will provide a canonical expression of roots in $\Phi_i(\mathscr{C})$ in terms of Π_i via π_i. However, since $\mathscr{C}$ lacks the integrality condition assumed in [21], the proof used in [21] will not apply here. Further, since it is no longer true that there exists a bijection between $\Phi_i(\mathscr{C})^+$ $(i = 1, 2)$ and the reflections in W, the proofs used in [11] and [16] will not apply either.

Let V_1' be a vector space over $\mathbb{R}$ with basis $\Pi_1' = \{\, \alpha_s' \mid s \in S \,\}$ in bijective correspondence with S, and let V_2' be a vector space over $\mathbb{R}$ with basis $\Pi_2' = \{\, \beta_s' \mid s \in S \,\}$, also in bijective correspondence with S. Define linear maps $\pi_1 \colon V_1' \to V_1$ and $\pi_2 \colon V_2' \to V_2$ by requiring that

$$\pi_1\Big(\sum_{s\in S} \lambda_s \alpha_s'\Big) = \sum_{s\in S} \lambda_s \alpha_s, \quad \text{and} \quad \pi_2\Big(\sum_{s\in S} \mu_s \beta_s'\Big) = \sum_{s\in S} \mu_s \beta_s,$$

for all $\lambda_s, \mu_s \in \mathbb{R}$, and define a bilinear map $\langle\, ,\, \rangle' \colon V_1' \times V_2' \to \mathbb{R}$ by requiring that $\langle \alpha_s', \beta_t' \rangle' = \langle \alpha_s, \beta_t \rangle$ for all $s, t \in S$. Observe that then $\langle x', y' \rangle' = \langle \pi_1(x'), \pi_2(y') \rangle$ for all $x' \in V_1'$ and $y' \in V_2'$.

Thus $\mathscr{C}' = (\, S, V_1', V_2', \Pi_1', \Pi_2', \langle\ ,\ \rangle' \,)$ is a free Coxeter datum with the same parameters as $\mathscr{C}$, and therefore is associated to the same abstract Coxeter system (W, R). Applying Theorems 2.10 to $\mathscr{C}'$ then yields isomorphisms $f_1' \colon W \to W_1'(\mathscr{C}')$ and $f_2' \colon W \to W_2'(\mathscr{C}')$. These isomorphisms induce W-actions on V_1' and V_2' via: $wx' = (f_1'(w))(x')$, and $uy' = (f_2'(u))(y')$ for all $w, u \in W$, $x' \in V_1'$ and $y' \in V_2'$. Note that for each $i \in \{1, 2\}$ and each $s \in S$, it follows from the above definitions that $f_i'(r_s) = \rho_{V_i'}(s)$, and this in turn yields that $\pi_i f_i'(r_s) = f_i(r_s)\pi_i$, where f_1 and f_2 are as in Theorem 2.10. Since W is generated by $\{\, r_s \mid s \in S \,\}$, it follows that $\pi_i f_i'(w) = f_i(w)\pi_i$ for all $w \in W$ and $i \in \{1, 2\}$. Summing up, one has that

$$\pi_i(wz') = w\pi_i(z') \qquad \text{for all } w \in W \text{ and } z' \in V_i', \tag{3.1}$$

and hence each π_i is a W-module homomorphism.

Proposition 3.5. *For each $i \in \{1, 2\}$, the restriction of π_i defines a W-equivariant bijection $\Phi_i(\mathscr{C}') \to \Phi_i(\mathscr{C})$.*

To prove Proposition 3.5, we need a few elementary results and some further notations.

Definition 3.6. For each $i \in \{1,2\}$, define an equivalence relation $\sim_i$ on $\Phi_i(\mathscr{C})$ as follows: if z_1 and $z_2 \in \Phi_i(\mathscr{C})$, then $z_1 \sim_i z_2$ if and only if z_1 and z_2 are (nonzero) scalar multiples of each other. For each $z \in \Phi_i(\mathscr{C})$, write $\widehat{z}$ for the equivalence class containing z, and set $\widehat{\Phi_i(\mathscr{C})} = \{\, \widehat{z} \mid z \in \Phi_i(\mathscr{C}) \,\}$.

Observe that the action of W on $\Phi_i(\mathscr{C})$ (for $i = 1, 2$) gives rise to a well-defined action of W on $\widehat{\Phi_i(\mathscr{C})}$ satisfying $w\widehat{z} = \widehat{wz}$ for all $w \in W$, and all $z \in \Phi_i(\mathscr{C})$.

Definition 3.7. For $i \in \{1,2\}$, and for each $w \in W$, define

$$N_i(w) = \{\, \widehat{\gamma} \mid \gamma \in \Phi_i^+(\mathscr{C}) \text{ and } w\gamma \in \Phi_i^-(\mathscr{C}) \,\}.$$

Note that for $w \in W$, the set $N_i(w)$ $(i = 1,2)$ can be alternatively characterized as $\{\, \widehat{\gamma} \mid \gamma \in \Phi_i^-(\mathscr{C}) \text{ and } w\gamma \in \Phi_i^+(\mathscr{C}) \,\}$. Hence $\widehat{z} \in N_i(w)$ if and only if precisely one element of the set $\{z, wz\}$ is in $\Phi_i^+(\mathscr{C})$. A mild generalization of the techniques used in ([17, 5.6 Proposition]) then yields the following result:

Lemma 3.8. (i) *If $s \in S$ then $N_1(r_s) = \{\widehat{\alpha_s}\}$ and $N_2(r_s) = \{\widehat{\beta_s}\}$.*
(ii) *Let $w \in W$. Then $N_1(w)$ and $N_2(w)$ both have cardinality $\ell(w)$.*
(iii) *Let w_1, $w_2 \in W$ and let $\dotplus$ denote set symmetric difference. Then $N_i(w_1w_2) = w_2^{-1}N_i(w_1) \dotplus N_i(w_2)$ for each $i \in \{1,2\}$.*
(iv) *Let w_1, $w_2 \in W$. Then for each $i \in \{1,2\}$,*

$$\ell(w_1w_2) = \ell(w_1) + \ell(w_2) \text{ if and only if } N_i(w_2) \subseteq N_i(w_1w_2).$$

□

If $w_1\,, w_2 \in W$ with $\ell(w_1w_2) = \ell(w_1) + \ell(w_2)$ then we call w_2 a *right hand segment* of w_1w_2. If $w = r_{s_1}r_{s_2}\cdots r_{s_l}$ (where $w \in W$ and $s_1, \cdots, s_l \in S$) with $\ell(w) = l$, then the above lemma yields that

$$N_1(w) = \{\widehat{\alpha_{s_l}},\, r_{s_l}\widehat{\alpha_{s_{l-1}}},\, r_{s_l}r_{s_{l-1}}\widehat{\alpha_{s_{l-2}}},\, \ldots,\, r_{s_l}r_{s_{l-1}}\ldots r_{s_2}\widehat{\alpha_{s_1}}\} \tag{3.2}$$

and

$$N_2(w) = \{\widehat{\beta_{s_l}},\, r_{s_l}\widehat{\beta_{s_{l-1}}},\, r_{s_l}r_{s_{l-1}}\widehat{\beta_{s_{l-2}}},\, ,\ldots,\, r_{s_l}r_{s_{l-1}}\ldots r_{s_2}\widehat{\beta_{s_1}}\}. \tag{3.3}$$

Essentially the same argument as in [15, Lemma, Lecture 3] yields the following:

Lemma 3.9. *W is finite if and only if $\widehat{\Phi_i(\mathscr{C})}$ is finite (for $i = 1,2$).*

□

Remark 3.10. Let $K \subseteq S$. If we define V_{1K} to be the subspace of V_1 spanned by $\Pi_{1K} = \{\, \alpha_s \mid s \in K \,\}$ and V_{2K} to be the subspace of V_2 spanned by $\Pi_{2K} = \{\, \beta_s \mid s \in K \,\}$, and let $\langle\ ,\ \rangle_K$ be the restriction of $\langle\ ,\ \rangle$ to $V_{1K} \times V_{2K}$, then clearly $\mathscr{C}_K = (K, V_{1K}, V_{2K}, \Pi_{1K}, \Pi_{2K}, \langle\ ,\ \rangle_K)$ is a Coxeter datum with parameters $\{\, m_{st} \mid s,t \in K \,\}$. Next we write $W_K = \langle \{\, r'_s \mid s \in K \,\} \rangle$ for the corresponding abstract Coxeter group, and let $\eta\colon W_K \to W$ be the homomorphism defined by $r'_s \mapsto r_s$ for all

$s \in K$. It follows immediately from the formulas for the actions of W on V_1 and W_K on V_{1K} that $r'_s v = r_s v$ for all $s \in K$ and $v \in V_{1K}$, and therefore $wv = \eta(w)v$ for all $w \in W_K$ and $v \in V_{1K}$. Since the action of W_K on V_{1K} is faithful, it follows that η is injective. Thus W_K can be identified with the *standard parabolic subgroup* of W generated by the set $\{\, r_s \mid s \in K \,\}$.

Definition 3.11. Given $K \subseteq S$, define $\Phi_{1K} (\subseteq V_{1K})$ and $\Phi_{2K} (\subseteq V_{2K})$ to be the root systems for W_K corresponding to $\mathscr{C}_K$, and Φ^+_{1K}, Φ^+_{2K} to be the corresponding sets of positive roots.

In other words, we have $\Phi_{1K} = \{\, w\alpha_r \mid w \in W_K \text{ and } r \in K \,\}$, and $\Phi^+_{1K} = \Phi_{1K} \cap \mathrm{PLC}(\Pi_{1K})$; similarly for Φ_{2K} and Φ^+_{2K}.

Remark 3.12. It is a particular case of Lemma 3.9 that W_K is finite if and only if either $\widehat{\Phi_{1K}}$ or $\widehat{\Phi_{2K}}$ is finite.

Remark 3.13. It follows easily from Lemma 3.2 (ii) that if W_K is finite then there exists a (unique) longest element $w_K \in W_K$ satisfying the condition $N_1(w_K) = \{\, \widehat{\alpha} \mid \alpha \in \Phi_{1K} \,\} = \widehat{\Phi_{1K}}$. For present purposes we require this only in the special case when K has cardinality 2.

Notation 3.14. *For r, $s \in S$, if $m_{rs} < \infty$ then $\langle\{r_r, r_s\}\rangle$ is finite, and its longest element, denoted by $w_{\{r,s\}}$, is $r_r r_s r_r \cdots = r_s r_r r_s \cdots$, where there are m_{rs} alternating factors on each side.*

Lemma 3.15. *Suppose that $s, t \in S$ and $w \in W$ such that $w\alpha_s = \lambda\alpha_t$ for some positive constant λ. Let $r \in S$ be such that $\ell(wr_r) < \ell(w)$. Then*

(i) *$\langle\{r_r, r_s\}\rangle$ is finite.*
(ii) *$N_1(w_{\{r,s\}} r_s) = \widehat{\Phi_{1\{\,r,s\,\}}} \setminus \{\widehat{\alpha_s}\}$.*
(iii) *$\ell(wr_s w^{-1}_{\{r,s\}}) = \ell(w) - \ell(w_{\{r,s\}} r_s)$.*

Proof. (i) Since $\ell(wr_r) < \ell(w)$, it follows from Lemma 3.2 (ii) that $w\alpha_r \in \Phi^-_1(\mathscr{C})$. Furthermore, $w\widehat{\alpha_r} \neq \widehat{\alpha_t}$, for otherwise $\widehat{\alpha_r} = \widehat{\alpha_s}$, contradicting Lemma 2.5. Observe that then

$$wr_s\alpha_r = w(\alpha_r - 2\langle\alpha_r, \beta_s\rangle\alpha_s) = \underbrace{w\alpha_r}_{\in\, \Phi^-_1(\mathscr{C}) \setminus \mathbb{R}\{\alpha_t\}} - \underbrace{2\lambda\langle\alpha_r, \beta_s\rangle\alpha_t}_{\text{a scalar multiple of } \alpha_t} .$$

Thus it can be checked that $wr_s\alpha_r \in \Phi^-_1(\mathscr{C})$. Now the fact that both $wr_s\alpha_r$ and $wr_s\alpha_s$ are negative implies that $wr_s(\lambda'\alpha_r + \mu'\alpha_s)$ is a negative linear combination of Π_1 whenever λ', $\mu' \geq 0$. This says precisely that $\widehat{\Phi_{1\{\,r,s\,\}}} \subseteq N_1(wr_s)$. Since $N_1(wr_s)$ is a finite set of size $\ell(wr_s)$ by Lemma 3.8 (ii), it follows from Remark 3.12 above that $\langle\{r_r, r_s\}\rangle$ must be finite.

(ii) First, $w_{\{r,s\}}$ exists by part (i). Next let $\mu\alpha_r + \nu\alpha_s \in \Phi^+_1(\mathscr{C})$ (where $\mu, \nu \geq 0$) be arbitrary. Then it follows from Remark 3.13 that $w_{\{r,s\}} r_s(\mu\alpha_r + \nu\alpha_s) \in \Phi^+_1(\mathscr{C})$ if and only if $r_s(\mu\alpha_r + \nu\alpha_s) \in \Phi^-_1(\mathscr{C})$, which by Lemma 3.8 (i) happens if and only if $\mu = 0$, and consequently $N_1(w_{\{r,s\}}r_s) = \widehat{\Phi_{1\,\{r,s\}}} \setminus \{\widehat{\alpha_s}\}$, as required.

(iii) By Lemma 3.8 (iv), to show that $w_{\{r,s\}}r_s$ is a right hand segment of w, it is enough to show that $N_1(w_{\{r,s\}}r_s) \subseteq N_1(w)$. Suppose that $\alpha \in \Phi_1^+(\mathscr{C})$ such that $\widehat{\alpha} \in N_1(w_{\{r,s\}}r_s)$. By (ii) above we may write $\alpha = c_1\alpha_r + c_2\alpha_s$ for some $c_1 > 0$ and $c_2 \geq 0$. Then $w\alpha = c_1 w\alpha_r + c_2\lambda\alpha_t$. Suppose for a contradiction that $w\alpha \in \Phi_1^+(\mathscr{C})$. Then $c_2\lambda\alpha_t = w\alpha - c_1 w\alpha_r \in \mathrm{PLC}(\Pi_1)$. Rearranging this equation gives $\lambda'\alpha_t = \sum_{t' \in S\setminus\{t\}} \lambda_{t'}\alpha_{t'}$, where λ' is a constant and $\lambda_{t'} \geq 0$ for all $t' \in S \setminus \{t\}$. Now if $\lambda' > 0$ then we have a contradiction to Lemma 2.5; on the other hand if $\lambda' \leq 0$ then we have a contradiction to $0 \notin \mathrm{PLC}(\Pi_1)$. ☐

Now we are ready to prove Proposition 3.5:

Proof of Proposition 3.5. Since $\Phi_1(\mathscr{C}') = \{\, w\alpha'_s \mid w \in W \text{ and } s \in S\,\}$, to prove that the restriction of π_1 to $\Phi_1(\mathscr{C}')$ is bijective it suffices to show that if $\pi_1(w\alpha'_s) = \pi_1(v\alpha'_t)$ for some $w, v \in W$ and $s, t \in S$, then $w\alpha'_s = v\alpha'_t$. Observe that $\pi_1(w\alpha'_s) = w\alpha_s$, and $\pi_1(v\alpha'_t) = v\alpha_t$. Hence it suffices to prove the following statement: if $w\alpha_s = \lambda\alpha_t$ for some $w \in W$, $s, t \in S$, and $\lambda \neq 0$, then $w\alpha'_s = \lambda\alpha'_t$. We assume that $w\alpha_s = \lambda\alpha_t$, and proceed by an induction on $\ell(w)$. The case $\ell(w) = 0$ reduces to the statement: if $\alpha_s = \lambda\alpha_t$ for some $s, t \in S$ then $\alpha'_s = \lambda\alpha'_t$. Given $\alpha_s = \lambda\alpha_t$, Lemma 2.5 and the requirement that $0 \notin \mathrm{PLC}(\Pi_1)$ together yield that $\lambda = 1$ and $s = t$, and we are done. Thus we may assume that $\ell(w) > 0$, and choose $r \in S$ such that $\ell(wr_r) < \ell(w)$. Lemma 3.15 yields that $\langle\{r_r, r_s\}\rangle$ is a finite dihedral group (hence m_{rs} is finite), and $\ell(w(w_{\{r,s\}}r_s)^{-1}) = \ell(w) - \ell(w_{\{r,s\}}r_s)$. We treat separately the cases m_{rs} even and m_{rs} odd.

If $m_{rs} = 2k$ is even, then $w_{\{r,s\}} = (r_r r_s)^k = (r_s r_r)^k$, and then the formulas in Lemma 2.6 (i) yield

$$\begin{aligned}(w_{\{r,s\}}r_s)\alpha_s &= -w_{\{r,s\}}\alpha_s = -(r_s r_r)^k\alpha_s \\ &= -\frac{\sin((m_{rs}+1)\pi/m_{rs})}{\sin(\pi/m_{rs})}\alpha_s - \frac{-\cos(\pi/m_{rs})}{\langle\alpha_r, \beta_s\rangle}\frac{\sin(\pi)}{\sin(\pi/m_{rs})}\alpha_r \\ &= \alpha_s; \qquad (3.4)\end{aligned}$$

and by exactly the same calculation in V_1' we have:

$$(w_{\{r,s\}}r_s)\alpha'_s = \alpha'_s. \qquad (3.5)$$

Observe that (3.4) yields that

$$\alpha_t = w\alpha_s = w(w_{\{r,s\}}r_s)^{-1}(w_{\{r,s\}}r_s)\alpha_s = w(w_{\{r,s\}}r_s)^{-1}\alpha_s. \qquad (3.6)$$

Now since $\ell(w(w_{\{r,s\}}r_s)^{-1}) < \ell(w)$, the inductive hypothesis combined with (3.6) give us

$$\alpha'_t = w(w_{\{r,s\}}r_s)^{-1}\alpha'_s. \qquad (3.7)$$

Then it follows from (3.5) and (3.7) that

$$w\alpha'_s = (w(w_{\{r,s\}}r_s)^{-1}(w_{\{r,s\}}r_s))\alpha'_s = w(w_{\{r,s\}}r_s)^{-1}\alpha'_s = \alpha'_t,$$

and the desired result follows by induction.

Next if $m_{rs} = 2k+1$ is odd, then $w_{\{r,s\}}r_s = \underbrace{\dots r_r r_s r_r}_{(m_{rs}-1)\text{ factors}} = (r_s r_r)^k$.

Then the formulas in Lemma 2.6 (i) yield that

$$\begin{aligned}(w_{\{r,s\}}r_s)\alpha_s &= (r_s r_r)^k \alpha_s\\ &= \frac{\sin(\pi)}{\sin(\pi/m_{rs})}\alpha_s + \frac{-\cos(\pi/m_{rs})}{\langle \alpha_r, \beta_s\rangle}\frac{\sin(2k\pi/m_{rs})}{\sin(\pi/m_{rs})}\alpha_r\\ &= \frac{-\cos(\pi/m_{rs})}{\langle \alpha_r, \beta_s\rangle}\alpha_r; \qquad (3.8)\end{aligned}$$

and by exactly the same calculation in V'_1 we have:

$$(w_{\{r,s\}}r_s)\alpha'_s = \frac{-\cos(\pi/m_{rs})}{\langle \alpha_r, \beta_s\rangle}\alpha'_r. \qquad (3.9)$$

Observe that (3.8) yields that

$$\alpha_t = w(w_{\{r,s\}}r_s)^{-1}(w_{\{r,s\}}r_s)\alpha_s = w(w_{\{r,s\}}r_s)^{-1}(\frac{-\cos(\pi/m_{rs})}{\langle \alpha_r, \beta_s\rangle}\alpha_r). \qquad (3.10)$$

Since $\ell(w(w_{\{r,s\}}r_s)^{-1}) < \ell(w)$, combining (3.10) and the inductive hypothesis yield that

$$\alpha'_t = w(w_{\{r,s\}}r_s)^{-1}(\frac{-\cos(\pi/m_{rs})}{\langle \alpha_r, \beta_s\rangle}\alpha'_r). \qquad (3.11)$$

Finally it follows from (3.9) and (3.11) that

$$\begin{aligned}w\alpha'_s = w(w_{\{r,s\}}r_s)^{-1}(w_{\{r,s\}}r_s)\alpha'_s &= w(w_{\{r,s\}}r_s)^{-1}(\frac{-\cos(\pi/m_{rs})}{\langle \alpha_r, \beta_s\rangle}\alpha'_r)\\ &= \alpha'_t,\end{aligned}$$

completing the proof that π_1 restricts to a bijection $\Phi_1(\mathscr{C}') \leftrightarrow \Phi_1(\mathscr{C})$. Exactly the same reasoning also yields that π_2 restricts to a bijection $\Phi_2(\mathscr{C}') \leftrightarrow \Phi_2(\mathscr{C})$. □

Under the freeness condition of $\mathscr{C}'$, each root $\alpha' \in \Phi_1(\mathscr{C}')$ can be written uniquely in the form $\alpha' = \sum_{s\in S}\lambda_s\alpha'_s$; we say that λ_s is the *coefficient* of α'_s in α', and denote it by $\mathrm{coeff}_s(\alpha')$. Similarly each root $\beta' \in \Phi_2(\mathscr{C}')$ can be written uniquely as $\beta' = \sum_{s\in S}\mu_s\beta'_s$; we say that μ_s is the *coefficient* of β'_s in β', and denote it by $\mathrm{coeff}_s(\beta')$.

Definition 3.16. (i) Let $\alpha \in \Phi_1(\mathscr{C})$ be arbitrary. For each $s \in S$, define the *canonical coefficient* of α_s in the root α, written $\mathrm{coeff}_s(\alpha)$, by requiring that $\mathrm{coeff}_s(\alpha) = \mathrm{coeff}_s(\pi_1^{-1}(\alpha))$, where π_1 is as in Proposition 3.5. The *support* of the root α, written $\mathrm{supp}(\alpha)$, is the set $\{\, \alpha_s \mid \mathrm{coeff}_s(\alpha) \neq 0 \,\}$.

(ii) Let $\beta \in \Phi_2(\mathscr{C})$ be arbitrary. For each $s \in S$, define the *canonical coefficient* of β_s in the root β, written $\operatorname{coeff}_s(\beta)$, by requiring that $\operatorname{coeff}_s(\beta) = \operatorname{coeff}_s(\pi_2^{-1}(\beta))$, where π_2 is as in Proposition 3.5. The *support* of the root β, written $\operatorname{supp}(\beta)$, is the set $\{ \beta_s \mid \operatorname{coeff}_s(\beta) \neq 0 \}$.

A similar proof as that of Proposition 3.5 establishes the following:

Proposition 3.17. *Suppose that $s, t \in S$. If $w\alpha_s = \lambda\alpha_t$ for some $w \in W$ and some non-zero constant λ, then $w\beta_s = \frac{1}{\lambda}\beta_t$.* □

In particular, Proposition 3.17 implies the following: if $w_1\alpha_s = w_2\alpha_t$ for some $s, t \in S$ and $w_1, w_2 \in W$, then $w_1\beta_s = w_2\beta_t$. Thus there exists a well-defined map $\Phi_1(\mathscr{C}) \to \Phi_2(\mathscr{C})$ satisfying the requirement that $w\alpha_s \mapsto w\beta_s$ for all $s \in S$ and $w \in W$. This is clearly the unique W-equivariant map $\Phi_1(\mathscr{C}) \to \Phi_2(\mathscr{C})$ satisfying $\alpha_s \mapsto \beta_s$ for all $s \in S$. Furthermore, by going through a similar reasoning as in the proof of Proposition 3.5, we may conclude that this map is a bijection.

Definition 3.18. Let $\phi : \Phi_1(\mathscr{C}) \to \Phi_2(\mathscr{C})$ be the W-equivariant bijection satisfying $\phi(\alpha_s) = \beta_s$ for all $s \in S$.

Remark 3.3 then immediately implies the following:

Corollary 3.19. *Let $\alpha \in \Phi_1(\mathscr{C})$. Then $\alpha \in \Phi_1^+(\mathscr{C})$ if and only if $\phi(\alpha) \in \Phi_2^+(\mathscr{C})$; and similarly $\alpha \in \Phi_1^-(\mathscr{C})$ if and only if $\phi(\alpha) \in \Phi_2^-(\mathscr{C})$.* □

Note that Proposition 3.17 can be generalized to the following:

Lemma 3.20. *Suppose that $\alpha \in \Phi_1(\mathscr{C})$, and suppose that λ is a non-zero constant such that $\lambda\alpha \in \Phi_1(\mathscr{C})$. Then $\phi(\lambda\alpha) = \frac{1}{\lambda}\phi(\alpha)$.*

Proof. Write $\alpha = w\alpha_s$ for some $w \in W$ and $s \in S$. The fact that $\lambda\alpha \in \Phi_1(\mathscr{C})$ implies that $w^{-1}(\lambda\alpha) = \lambda w^{-1}\alpha = \lambda\alpha_s \in \Phi_1(\mathscr{C})$. Then $\phi(\lambda\alpha_s) = \frac{1}{\lambda}\phi(\alpha_s)$ by Proposition 3.17. Now it follows from the W-equivariance of ϕ that

$$\phi(\lambda\alpha) = \phi(\lambda w\alpha_s) = \phi(w\lambda\alpha_s) = w\phi(\lambda\alpha_s) = \frac{1}{\lambda}w\phi(\alpha_s) = \frac{1}{\lambda}\phi(\alpha).$$

□

Definition 3.21. Let $\mathscr{C} = (\, S, V_1, V_2, \Pi_1, \Pi_2, \langle\ ,\ \rangle\,)$ be a Coxeter datum.
(i) For each $i \in \{1, 2\}$ and each $z \in \Phi_i^+(\mathscr{C})$, define the *depth* of z (written $\operatorname{dp}_{\mathscr{C},i}(z)$) to be

$$\operatorname{dp}_{\mathscr{C},i}(z) = \min\{\, \ell(w) \mid w \in W \text{ and } wz \in \Phi_i^-(\mathscr{C}) \,\}.$$

(ii) For each $i \in \{1, 2\}$ and $z_1, z_2 \in \Phi_i^+(\mathscr{C})$, write $z_1 \preceq_i z_2$ if there exists $w \in W$ such that $z_2 = wz_1$, and $\operatorname{dp}_{\mathscr{C},i}(z_2) = \operatorname{dp}_{\mathscr{C},i}(z_1) + \ell(w)$. Furthermore, we write $z_1 \prec_i z_2$ if $z_1 \preceq_i z_2$ but $z_1 \neq z_2$.

A mild generalization of Lemma 1.7 of [3] yields the following:

Lemma 3.22. *Suppose that* $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\,,\,\rangle\,)$ *is a Coxeter datum. Let* $s \in S$, $\alpha \in \Phi_1^+(\mathscr{C}) \setminus \mathbb{R}\{\alpha_s\}$, *and* $\beta \in \Phi_2^+(\mathscr{C}) \setminus \mathbb{R}\{\beta_s\}$. *Then*

$$\mathrm{dp}_{\mathscr{C},1}(r_s\alpha) = \begin{cases} \mathrm{dp}_{\mathscr{C},1}(\alpha) - 1 & \text{if } \langle \alpha, \beta_s \rangle > 0, \\ \mathrm{dp}_{\mathscr{C},1}(\alpha) & \text{if } \langle \alpha, \beta_s \rangle = 0, \\ \mathrm{dp}_{\mathscr{C},1}(\alpha) + 1 & \text{if } \langle \alpha, \beta_s \rangle < 0; \end{cases}$$

and

$$\mathrm{dp}_{\mathscr{C},2}(r_s\beta) = \begin{cases} \mathrm{dp}_{\mathscr{C},2}(\beta) - 1 & \text{if } \langle \alpha_s, \beta \rangle > 0, \\ \mathrm{dp}_{\mathscr{C},2}(\beta) & \text{if } \langle \alpha_s, \beta \rangle = 0, \\ \mathrm{dp}_{\mathscr{C},2}(\beta) + 1 & \text{if } \langle \alpha_s, \beta \rangle < 0. \end{cases}$$

□

Lemma 3.23. *Suppose that* $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\,,\,\rangle\,)$ *is a Coxeter datum, and* $\alpha \in \Phi_1^+(\mathscr{C})$. *Then* $\mathrm{dp}_{\mathscr{C},1}(\alpha) = \mathrm{dp}_{\mathscr{C},2}(\phi(\alpha))$.

Proof. Let $w \in W$ be such that $w\alpha \in \Phi_1^-(\mathscr{C})$ and $\mathrm{dp}_{\mathscr{C},1}(\alpha) = \ell(w)$. Then Corollary 3.19 yields that $\phi(w\alpha) \in \Phi_2^-(\mathscr{C})$, and it follows from the W-equivariance of ϕ that, $w(\phi(\alpha)) \in \Phi_2^-(\mathscr{C})$. Hence

$$\mathrm{dp}_{\mathscr{C},2}(\phi(\alpha)) \leq \ell(w) = \mathrm{dp}_{\mathscr{C},1}(\alpha).$$

By symmetry, $\mathrm{dp}_{\mathscr{C},1}(\alpha) \leq \mathrm{dp}_{\mathscr{C},2}(\phi(\alpha))$, whence equality. □

Lemma 3.24. *Suppose that* $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\,,\,\rangle\,)$ *is a Coxeter datum. Let* $\alpha \in \Phi_1(\mathscr{C})$ *and* $s \in S$. *Then*

$$\langle \alpha, \beta_s \rangle > 0 \text{ if and only if } \langle \alpha_s, \phi(\alpha) \rangle > 0;$$

and

$$\langle \alpha, \beta_s \rangle = 0 \text{ if and only if } \langle \alpha_s, \phi(\alpha) \rangle = 0;$$

and

$$\langle \alpha, \beta_s \rangle < 0 \text{ if and only if } \langle \alpha_s, \phi(\alpha) \rangle < 0.$$

Proof. Combine Lemma 3.22 and Lemma 3.23, then the desired result follows. □

In fact, the above gives rise to a more general result:

Corollary 3.25. *Suppose that* $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\,,\,\rangle\,)$ *is a Coxeter datum, and* $\alpha_1, \alpha_2 \in \Phi_1(\mathscr{C})$. *Then*

$$\langle \alpha_1, \phi(\alpha_2) \rangle > 0 \quad \text{if and only if} \quad \langle \alpha_2, \phi(\alpha_1) \rangle > 0;$$

and

$$\langle \alpha_1, \phi(\alpha_2) \rangle = 0 \quad \text{if and only if} \quad \langle \alpha_2, \phi(\alpha_1) \rangle = 0;$$

and

$$\langle \alpha_1, \phi(\alpha_2) \rangle < 0 \quad \text{if and only if} \quad \langle \alpha_2, \phi(\alpha_1) \rangle < 0.$$

Proof. Write $\alpha_2 = w\alpha_s$ for some $w \in W$ and $s \in S$. Given the W-invariance of $\langle\,,\,\rangle$ and the W-equivariance of ϕ, we have

$$\langle \alpha_1, \phi(\alpha_2) \rangle = \langle \alpha_1, w\beta_s \rangle = \langle w^{-1}\alpha_1, \beta_s \rangle.$$

It then follows from Lemma 3.24 that $\langle w^{-1}\alpha_1, \beta_s\rangle > 0$ precisely when

$$\langle \alpha_s, \phi(w^{-1}\alpha_1)\rangle = \langle w\alpha_s, \phi(\alpha_1)\rangle = \langle \alpha_2, \phi(\alpha_1)\rangle > 0.$$

The rest of the desired result follows in a similar way. □

4. Comparison with the Standard Geometric Realization of Coxeter Groups

In this section we recover the root systems of Coxeter groups in the sense of [13] or [17] as special cases of root systems arising from a Coxeter datum. Subsequently we give comparison results between such special cases and the more general root systems arising from a Coxeter datum. These comparisons will provide useful reduction of the non-orthogonal representations studied in the previous sections into those of [13] or [17].

Fix a Coxeter datum $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\,,\,\rangle\,)$, and let V be a vector space over $\mathbb{R}$ with a basis $\Pi = \{\,\gamma_s \mid s \in S\,\}$ in bijective correspondence with S. Suppose that $(\,,\,)\colon V \times V \to \mathbb{R}$ is a symmetric bilinear form satisfying the following conditions:

(C1') $(\gamma_s, \gamma_s) = 1$, for all $s \in S$;
(C2') $(\gamma_s, \gamma_t) \leq 0$, for all distinct $s, t \in S$;
(C3') $(\gamma_s, \gamma_t)^2 = \langle \alpha_s, \beta_t\rangle\langle \alpha_t, \beta_s\rangle$, for all $s, t \in S$.

Then $\mathscr{C}'' = (\,S, V, V, \Pi, \Pi, (\,,\,)\,)$ is a free Coxeter datum with the same Coxeter parameters m_{st}, $(s, t \in S)$ as those of $\mathscr{C}$, and hence $\mathscr{C}$ and $\mathscr{C}''$ are associated to the same abstract Coxeter system (W, R). It is easily checked that $\Phi_1(\mathscr{C}'') = \Phi_2(\mathscr{C}'')$, allowing us to write Φ in places of $\Phi_1(\mathscr{C}'')$ and $\Phi_2(\mathscr{C}'')$. Furthermore, we write Φ^+ and Φ^- for the corresponding set of positive roots and negative roots respectively. It is also readily checked that W can be faithfully embedded into the orthogonal group of the bilinear form $(\,,\,)$ on V. This recovers the orthogonal representation of W as defined in [2], [13] and [17]. We refer to such V as the *standard geometric realization (Tits representation)* of W or simply the *standard Tits representation* of W.

It follows from Lemma 2.13 that $(\,,\,)$ is W-invariant. It is well-known (and can be readily checked) that if x and λx are both in Φ for some constant λ, then $\lambda = \pm 1$. Since $\mathscr{C}''$ is free, it follows that each $\gamma \in \Phi$ can be written uniquely in the form $\gamma = \sum_{s \in S} \lambda_s \gamma_s$. We say that λ_s is the *coefficient* of γ_s in γ, and denote it by $\operatorname{coeff}_s(\gamma)$. Finally, to simplify notation, for any $\gamma \in \Phi$ we write $\operatorname{dp}(\gamma)$ in place of $\operatorname{dp}_{\mathscr{C}'',1}(\gamma)(= \operatorname{dp}_{\mathscr{C}'',2}(\gamma))$, and for $\gamma_1, \gamma_2 \in \Phi$ we simply write $\gamma_1 \preceq \gamma_2$ when $\gamma_1 \preceq_1 \gamma_2$ (and $\gamma_1 \preceq_2 \gamma_2$).

Similar arguments as for Proposition 3.5 gives us the following:

Proposition 4.1. *There are W-equivariant maps $\phi_1\colon \Phi_1(\mathscr{C}) \to \Phi$, and $\phi_2\colon \Phi_2(\mathscr{C}) \to \Phi$, satisfying $\phi_1(\alpha_s) = \gamma_s = \phi_2(\beta_s)$, for all $s \in S$.* □

Remark 4.2. We stress that unlike π_1, π_2 of Proposition 3.5 and ϕ of Definition 3.18, the new maps ϕ_1 and ϕ_2 are not injective in general, and we shall see more on this fact in Lemma 4.8 (ii) below.

Lemma 4.3. *Suppose that* $\alpha \in \Phi_1(\mathscr{C})$. *Then* $\alpha \in \Phi_1^+(\mathscr{C})$ *implies that* $\phi_1(\alpha) \in \Phi^+$, *and* $\alpha \in \Phi_1^-(\mathscr{C})$ *implies that* $\phi_1(\alpha) \in \Phi^-$.

Proof. We may write $\alpha = w\alpha_r$, for suitable $w \in W$ and $r \in S$. If $\alpha \in \Phi_1^+(\mathscr{C})$, then Lemma 3.2 (ii) yields that $\ell(wr_r) = \ell(w) + 1$, in which case Lemma 3.2 (ii) applied to the Coxeter datum $\mathscr{C}''$ yields that $\phi_1(\alpha) = \phi_1(w\alpha_r) = w\phi_1(\alpha_r) = w\gamma_r \in \Phi^+$. Likewise we see that $\alpha \in \Phi_1^-(\mathscr{C})$ implies that $\phi_1(\alpha) \in \Phi^-$. $\square$

Using the same argument as in Lemma 3.23 we have:

Lemma 4.4. *Suppose that* $\alpha \in \Phi_1^+(\mathscr{C})$. *Then* $\mathrm{dp}_{\mathscr{C},1}(\alpha) = \mathrm{dp}(\phi_1(\alpha))$. $\square$

Corollary 4.5. *Suppose that* $\alpha \in \Phi_1(\mathscr{C})$, *and* $s \in S$. *Then*

$$(\phi_1(\alpha), \gamma_s) > 0 \text{ if and only if } \langle \alpha, \beta_s \rangle > 0,$$

and

$$(\phi_1(\alpha), \gamma_s) = 0 \text{ if and only if } \langle \alpha, \beta_s \rangle = 0,$$

and

$$(\phi_1(\alpha), \gamma_s) < 0 \text{ if and only if } \langle \alpha, \beta_s \rangle < 0.$$

Proof. Follows from Lemma 4.4 and Lemma 3.22 applied to $\mathscr{C}''$. $\square$

Next we have a well-known result on Coxeter groups (a proof can be found in [5], in the discussion immediately before Lemma 2.1) which we shall use repeatedly in later calculations.

Lemma 4.6. *Suppose that* $I \subseteq S$ *and* $w \in W$. *Let* W_I *denote the standard parabolic subgroup in* W *corresponding to* I, *as defined in Remark 3.10. Choose* $w' \in wW_I$ *to be of minimal length in the left coset of* W_I *in* W *containing* w. *Then* $\ell(w'v) = \ell(w') + \ell(v)$ *for all* $v \in W_I$. $\square$

Proposition 4.7. *For each* $\alpha \in \Phi_1(\mathscr{C})$ *and each* $r \in S$,

$$\mathrm{coeff}_r(\alpha)\,\mathrm{coeff}_r(\phi(\alpha)) \geq (\mathrm{coeff}_r(\phi_1(\alpha)))^2.$$

Proof. Replace α by $-\alpha$ if needed, we may assume that $\alpha \in \Phi_1^+(\mathscr{C})$, and we may write $\alpha = w\alpha_s$, where $w \in W$ and $s \in S$. The proof is based on an induction on $\ell(w)$. If $\ell(w) = 0$ then the result clearly holds with an equality. Thus we may assume that $\ell(w) \geq 1$, and choose $t \in S$ such that $\ell(wr_t) = \ell(w) - 1$. Observe that $w = w_1w_2$, where w_1 is of minimal length in the coset $w\langle\{r_s, r_t\}\rangle$, and w_2 is an alternating product of r_s and r_t, ending in r_t. Then Lemma 4.6 yields that $\ell(w) = \ell(w_1) + \ell(w_2)$, $\ell(w_1r_s) = \ell(w_1) + 1$, and $\ell(w_1r_t) = \ell(w_1) + 1$. Consequently $w_1\alpha_s \in \Phi_1^+(\mathscr{C})$ and $w_1\alpha_t \in \Phi_1^+(\mathscr{C})$ by Lemma 3.2 (ii).

The formulas in Lemma 2.6 yield that $w_2\alpha_s = p\alpha_s + \lambda q\alpha_t$, where λ is a positive constant and $pq \geq 0$. If p and q are both negative, then

$$\alpha = w\alpha_s = w_1w_2\alpha_s = w_1(p\alpha_s + \lambda q\alpha_t) = pw_1\alpha_s + \lambda qw_1\alpha_t \in \Phi_1^-(\mathscr{C}),$$

contradicting the assumption that $\alpha \in \Phi_1^+(\mathscr{C})$. Therefore $p, q \geq 0$. By Lemma 2.5, $\{\alpha_s, \alpha_t\}$ is linearly independent, hence $\mathrm{coeff}_s(w_2\alpha_s) = p$, and $\mathrm{coeff}_t(w_2\alpha_s) = \lambda q$. Similar calculations as in Lemma 2.6 yield that

$$w_2\gamma_s = p\gamma_s + q\gamma_t, \quad \text{and} \quad w_2\beta_s = p\beta_s + \tfrac{q}{\lambda}\beta_t.$$

And Lemma 2.5 implies that $\mathrm{coeff}_s(w_2\beta_s) = p$ and $\mathrm{coeff}_t(w_2\beta_s) = \frac{q}{\lambda}$. Next we set

$$x = \mathrm{coeff}_r(\alpha)\ ,\ x' = \mathrm{coeff}_r(\phi(\alpha))\ ,\ x'' = \mathrm{coeff}_r(\phi_1(\alpha));$$

and

$$y = \mathrm{coeff}_r(w_1\alpha_s)\ ,\ y' = \mathrm{coeff}_r(w_1\beta_s)\ ,\ y'' = \mathrm{coeff}_r(w_1\gamma_s);$$

and

$$z = \mathrm{coeff}_r(w_1\alpha_t)\ ,\ z' = \mathrm{coeff}_r(w_1\beta_t)\ ,\ z'' = \mathrm{coeff}_r(w_1\gamma_t).$$

Then

$$\begin{aligned} xx' - x''^2 &= (py + \lambda qz)(py' + \tfrac{1}{\lambda}qz') - (py'' + qz'')^2 \\ &= p^2(yy' - y''^2) + q^2(zz' - z''^2) + pq(\tfrac{1}{\lambda}yz' + \lambda zy' - 2y''z''). \end{aligned} \tag{4.1}$$

Since $\ell(w_1) < \ell(w)$, it follows from the inductive hypothesis that

$$yy' \geq (y'')^2 \qquad \text{and} \qquad zz' \geq (z'')^2,$$

and so the first two summands in the last line of (4.1) are nonnegative. Next apply the geometric mean and arithmetic mean inequality to the terms $\frac{1}{\lambda}yz'$ and $\lambda y'z$, and we can conclude that $\frac{1}{\lambda}yz' + \lambda y'z \geq 2\sqrt{yy'zz'}$. But the inductive hypothesis yields that $yy'zz' \geq (y''z'')^2$, showing that the third summand in the last line of(4.1) is also nonnegative, whence $xx' - x''^2 \geq 0$, and the desired result follows by induction. ☐

Lemma 4.8. *Suppose that $\alpha \in \Phi_1(\mathscr{C})$.*
(i) Let $t \in S$. Then

$$\begin{cases} \mathrm{coeff}_t(\alpha) = 0 & \textit{if and only if } \mathrm{coeff}_t(\phi_1(\alpha)) = 0, \\ \mathrm{coeff}_t(\alpha) > 0 & \textit{if and only if } \mathrm{coeff}_t(\phi_1(\alpha)) > 0, \\ \mathrm{coeff}_t(\alpha) < 0 & \textit{if and only if } \mathrm{coeff}_t(\phi_1(\alpha)) < 0. \end{cases}$$

(ii) The situation $\phi_1(\alpha) = \pm\gamma_s$ $(s \in S)$ arises if and only if $\alpha = \lambda\alpha_s$, for some nonzero constant λ.

Proof. (i) Lemma 3.2 (i) applied to the Coxeter datum $\mathscr{C}''$ yields that $\Phi = \Phi^+ \uplus \Phi^-$. Hence we only need to verify that $\mathrm{coeff}_t(\alpha) = 0$ if and only if $\mathrm{coeff}_t(\phi_1(\alpha)) = 0$, for the rest of (i) follows readily from Lemma 4.3. By Proposition 4.7 we only need to show that the condition $\mathrm{coeff}_t(\phi_1(\alpha)) = 0$ implies that $\mathrm{coeff}_t(\alpha) = 0$. Replacing α by $-\alpha$ if needs be, we may assume that $\alpha \in \Phi_1^+(\mathscr{C})$, and write $\alpha = w\alpha_r$ for some $w \in W$ and $r \in S$. If $\ell(w) = 0$ then there is nothing to prove. Thus we may assume that $\ell(w) \geq 1$ and proceed by an induction on $\ell(w)$. Choose $s \in S$ such that $\ell(wr_s) = \ell(w) - 1$. Now $w = w_1 w_2$, where $w_1 \in w\langle\{r_r, r_s\}\rangle$ is of minimal length, and w_2 is an alternating product of r_r and r_s ending in r_s. Then Lemma 4.6 implies that w_2 is a right segment of w. Since $\alpha = w\alpha_r$ is positive, it follows from Lemma 3.8 (iv) that $w_2\alpha_r$ is positive too. Direct calculations as those in Lemma 2.6 show that

$$w_2\gamma_r = p\gamma_r + q\gamma_s, \quad w_2\alpha_r = p\alpha_r + \lambda q\alpha_s \text{ and } w_2\beta_r = p\beta_r + \tfrac{q}{\lambda}\beta_s$$

for some non-negative constants p and q, and some positive constant λ. Now Lemma 2.5 yields that $\operatorname{coeff}_r(w_2\alpha_r) = p$, $\operatorname{coeff}_s(w_2\alpha_r) = \lambda q$, $\operatorname{coeff}_r(w_2\beta_r) = p$ and $\operatorname{coeff}_s(w_2\beta_r) = \frac{q}{\lambda}$. Then

$$\begin{aligned} 0 = \operatorname{coeff}_t(\phi_1(\alpha)) &= \operatorname{coeff}_t(w\gamma_r) = \operatorname{coeff}_t(w_1(p\gamma_r + q\gamma_s)) \\ &= p\operatorname{coeff}_t(w_1\gamma_r) + q\operatorname{coeff}_t(w_1\gamma_s). \end{aligned}$$

By Lemma 4.6 and Lemma 3.2 (ii) (applied to the Coxeter datum $\mathscr{C}''$), $w_1\gamma_r$ and $w_1\gamma_s$ are both positive, and hence it follows from the above that

$$p\operatorname{coeff}_t(w_1\gamma_r) = q\operatorname{coeff}_t(w_1\gamma_s) = 0.$$

Then the inductive hypothesis yields that

$$p\operatorname{coeff}_t(w_1\alpha_r) = q\operatorname{coeff}_t(w_1\alpha_s) = 0,$$

and therefore

$$\operatorname{coeff}_t(\alpha) = \operatorname{coeff}_t(w\alpha_r) = p\operatorname{coeff}_t(w_1\alpha_r) + \lambda q\operatorname{coeff}_t(w_1\alpha_s) = 0$$

as required, and (i) follows by induction.
(ii) Follows readily from part (i) above. □

Remark 4.9. Suppose that $\beta \in \Phi_2(\mathscr{C})$, and $t \in S$. Then the same arguments as those used in the proof of the above Lemma yield that

$$\begin{cases} \operatorname{coeff}_t(\beta) = 0 & \text{if and only if } \operatorname{coeff}_t(\phi_2(\beta)) = 0, \\ \operatorname{coeff}_t(\beta) > 0 & \text{if and only if } \operatorname{coeff}_t(\phi_2(\beta)) > 0, \\ \operatorname{coeff}_t(\beta) < 0 & \text{if and only if } \operatorname{coeff}_t(\phi_2(\beta)) < 0. \end{cases}$$

The next result is taken from [4]:

Lemma 4.10. (Brink [4, Proposition 2.1]) *Suppose that* $\gamma \in \Phi$ *and* $r \in S$. *Then* $\operatorname{coeff}_r(\gamma) > 0$ *implies that* $\operatorname{coeff}_r(\gamma) \geq 1$.

Further, suppose that $0 < \operatorname{coeff}_r(\gamma) < 2$. *Then either* $\operatorname{coeff}_r(\gamma) = 1$ *or* $\operatorname{coeff}_r(\gamma) = 2\cos(\frac{\pi}{m_{r_1 r_2}})$, *where* r_1, $r_2 \in S$ *with* $4 \leq m_{r_1 r_2} < \infty$. □

Combining the results in Proposition 4.7 to Lemma 4.10 (inclusive) and an argument similar to the one used in the proof of Proposition 4.7, we may deduce the following:

Proposition 4.11. *Let $\alpha \in \Phi_1(\mathscr{C})$ and $t \in S$ be arbitrary. Then*
$$\operatorname{coeff}_t(\alpha) > 0 \text{ if and only if } \operatorname{coeff}_t(\phi(\alpha)) > 0,$$
and in this case $\operatorname{coeff}_t(\alpha)\operatorname{coeff}_t(\phi(\alpha)) \geq 1$. In particular,
$$\phi(\operatorname{supp}(\alpha)) = \operatorname{supp}(\phi(\alpha)).$$
Furthermore, suppose that $1 \leq \operatorname{coeff}_t(\alpha)\operatorname{coeff}_t(\phi(\alpha)) < 4$. Then either
$$\operatorname{coeff}_t(\alpha)\operatorname{coeff}_t(\phi(\alpha)) = 1, \text{ or else } \operatorname{coeff}_t(\alpha)\operatorname{coeff}_t(\phi(\alpha)) = 4\cos^2(\frac{\pi}{m}),$$
where $m = m_{r_1 r_2}$, for some $r_1, r_2 \in S$ with $4 \leq m < \infty$. In particular, if $\alpha \in \Phi_1^+(\mathscr{C})$ then $\operatorname{coeff}_t(\alpha)\operatorname{coeff}_t(\phi(\alpha)) = 1$ if and only if $\operatorname{coeff}_t(\phi_1(\alpha)) = 1$. □

Proposition 4.12. *Suppose that $\alpha_1, \alpha_2 \in \Phi_1(\mathscr{C})$. Then*
$$\langle \alpha_1, \phi(\alpha_2)\rangle\langle \alpha_2, \phi(\alpha_1)\rangle \geq (\phi_1(\alpha_1), \phi_1(\alpha_2))^2.$$

Proof. Since both $\langle\, ,\, \rangle$ and $(\, ,\,)$ are W-invariant, and ϕ, ϕ_1 are W-equivariant, we may replace α_1 and α_2 by $u\alpha_1$ and $u\alpha_2$ for a suitable $u \in W$ so that $\alpha_2 = \alpha_s$ for some $s \in S$. Furthermore, replace α_1 by $-\alpha_1$ if needs be, we may assume that $\alpha_1 \in \Phi_1^+(\mathscr{C})$. We proceed with an induction on the depth of α_1.

If $\operatorname{dp}_{\mathscr{C},1}(\alpha_1) = 1$, then $\alpha_1 = \lambda\alpha_r$, for some positive constant λ and some $r \in S$. It follows from Lemma 4.8 (ii) that $\phi_1(\alpha_1) = \gamma_r$. Furthermore, Lemma 3.20 yields that $\phi(\alpha_1) = \frac{1}{\lambda}\beta_r$, and hence
$$\begin{aligned}\langle \alpha_1, \phi(\alpha_2)\rangle\langle \alpha_2, \phi(\alpha_1)\rangle &= \lambda\langle \alpha_r, \beta_s\rangle\tfrac{1}{\lambda}\langle \alpha_s, \beta_r\rangle \\ &= \langle \alpha_r, \beta_s\rangle\langle \alpha_s, \beta_r\rangle = (\gamma_r, \gamma_s)^2 = (\phi_1(\alpha_1), \phi_1(\alpha_2)).\end{aligned}$$
Thus we may assume that $\operatorname{dp}_{\mathscr{C},1}(\alpha_1) > 1$. First, if $\langle \alpha_1, \beta_s\rangle > 0$ then Lemma 3.22 yields that $r_s\alpha_1 \prec_1 \alpha_1$, and hence
$$\begin{aligned}\langle \alpha_1, \beta_s\rangle\langle \alpha_s, \phi(\alpha_1)\rangle &= \langle r_s\alpha_1, r_s\beta_s\rangle\langle r_s\alpha_s, r_s\phi(\alpha_1)\rangle \\ &= (-\langle r_s\alpha_1, \beta_s\rangle)(-\langle \alpha_s, \phi(r_s\alpha_1)\rangle) \\ &\geq (\phi_1(r_s\alpha_1), \gamma_s)^2 \quad \text{(by the inductive hypothesis)} \\ &= (\phi_1(\alpha_1), -\gamma_s)^2 \\ &= (\phi_1(\alpha_1), \gamma_s)^2,\end{aligned}$$
as required. Thus we may further assume that $\langle \alpha_1, \beta_s\rangle \leq 0$.

Next let $t \in S$ be such that $r_t\alpha_1 \prec_1 \alpha_1$. Then Lemma 3.22 yields that $\langle \alpha_1, \beta_t\rangle > 0$, and, in particular, $s \neq t$. Now let $w \in W$ be a maximal length alternating product of r_s and r_t ending in r_t such that $\operatorname{dp}_{\mathscr{C},1}(w\alpha_1) = \operatorname{dp}_{\mathscr{C},1}(\alpha_1) - \ell(w)$. In particular, $w \neq 1$. Thus $\operatorname{dp}_{\mathscr{C},1}(w\alpha_1) \lneq \operatorname{dp}_{\mathscr{C},1}(\alpha_1)$, and so the inductive hypothesis yields that
$$\langle w\alpha_1, \beta_s\rangle\langle \alpha_s, \phi(w\alpha_1)\rangle \geq (\gamma_s, \phi_1(w\alpha_1))^2 \tag{4.2}$$
and
$$\langle w\alpha_1, \beta_t\rangle\langle \alpha_t, \phi(w\alpha_1)\rangle \geq (\gamma_t, \phi_1(w\alpha_1))^2. \tag{4.3}$$

Next, for any reduced expression $w = r_{s_1}\cdots r_{s_l}$ ($s_i \in S$), it is readily checked that $r_{s_l}\alpha_1 \prec_1 \alpha_1$, and so it follows from Lemma 3.22 that $\langle \alpha_1, \beta_{s_l}\rangle > 0$. Therefore $s_l \neq s$, and w has no reduced expression ending in r_s. That is, w is a product of r_s and r_t with strictly fewer than m_{st} factors. Thus by Lemma 2.11 or a direct calculation as that in Lemma 2.6, there are non-negative constants p, q and positive constant λ such that

$$w\gamma_s = p\gamma_s + q\gamma_t \quad\text{and}\quad w\alpha_s = p\alpha_s + \lambda q\alpha_t \quad\text{and}\quad w\beta_s = p\beta_s + \tfrac{q}{\lambda}\beta_t.$$

Thus

$$\begin{aligned}
&\langle \alpha_1, \beta_s\rangle\langle \alpha_s, \phi(\alpha_1)\rangle - (\gamma_s, \phi_1(\alpha_1))^2\\
&= \langle w\alpha_1, w\beta_s\rangle\langle w\alpha_s, \phi(w\alpha_1)\rangle - (w\gamma_s, \phi_1(w\alpha_1))^2\\
&\qquad\qquad\qquad\text{(since } \langle\ ,\ \rangle \text{ and } (\ ,\ ) \text{ are } W\text{-invariant)}\\
&= \langle w\alpha_1, p\beta_s + \tfrac{q}{\lambda}\beta_t\rangle\langle p\alpha_s + \lambda q\alpha_t, \phi(w\alpha_1)\rangle - (\phi_1(w\alpha_1), p\gamma_s + q\gamma_t)^2\\
&= \underbrace{p^2(\langle w\alpha_1, \beta_s\rangle\langle \alpha_s, \phi(w\alpha_1)\rangle - (\phi_1(w\alpha_1), \gamma_s)^2)}_{\text{A}}\\
&\qquad + \underbrace{q^2(\langle w\alpha_1, \beta_t\rangle\langle \alpha_t, \phi(w\alpha_1)\rangle - (\phi_1(w\alpha_1), \gamma_t)^2)}_{\text{B}} + C.
\end{aligned}$$

where

$$\begin{aligned}
C = pq(\ &\tfrac{1}{\lambda}\langle w\alpha_1, \beta_t\rangle\langle \alpha_s, \phi(w\alpha_1)\rangle + \lambda\langle w\alpha_1, \beta_s\rangle\langle \alpha_t, \phi(w\alpha_1)\rangle\\
&- 2(\phi_1(w\alpha_1), \gamma_s)(\phi_1(w\alpha_1), \gamma_t)\ ).
\end{aligned}$$

It follows from (4.2) and (4.3) that A and B are both nonnegative. It follows from the geometric mean and arithmetic mean inequality that

$$\begin{aligned}
&\tfrac{1}{\lambda}\langle w\alpha_1, \beta_t\rangle\langle \alpha_s, \phi(w\alpha_1)\rangle + \lambda\langle w\alpha_1, \beta_s\rangle\langle \alpha_t, \phi(w\alpha_1)\rangle\\
&\qquad \geq 2\sqrt{\langle w\alpha_1, \beta_s\rangle\langle \alpha_s, \phi(w\alpha_1)\rangle\langle w\alpha_1, \beta_t\rangle\langle \alpha_t, \phi(w\alpha_1)\rangle}\\
&\qquad \geq 2(\phi_1(w\alpha_1), \gamma_s)(\phi_1(w\alpha_1), \gamma_t) \qquad \text{(by (4.2) and (4.3))},
\end{aligned}$$

that is, $C \geq 0$ as well. Therefore $\langle \alpha_1, \beta_s\rangle\langle \alpha_s, \phi(\alpha_1)\rangle \geq (\gamma_s, \phi_1(\alpha_1))^2$, and the desired result follows by induction. □

5. TITS CONES AND A NON-POSITIVITY RESULT

Let $\mathscr{C} = (\,S, V_1, V_2, \Pi_1, \Pi_2, \langle\ ,\ \rangle\,)$ be a fixed Coxeter datum, and let (W, R) be the corresponding Coxeter system. In this section we study a class of cones associated to $\mathscr{C}$ that are analogous to the *Tits cones* in the classical setting (as defined in [22] or [17]). Furthermore, we investigate certain W-invariant sets in V_1 and V_2 that are closely related to these cones. The key result of this section is a generalization to Proposition 1.2 of [20] and Proposition 3.4 of [16]. For this section we impose one additional condition on $\mathscr{C}$, namely:

(C6) $V_1 = \operatorname{span}(\Pi_1)$ and $V_2 = \operatorname{span}(\Pi_2)$,

and we retain all other conventions and notations of earlier sections.

Notation 5.1. *For each* $i \in \{1,2\}$ *and* $I \subseteq S$*, recall the notations* Π_{iI} *and* V_{iI} *introduced in Remark 3.10, and set* $P_{iI} = \mathrm{PLC}(\Pi_{iI}) \cup \{0\}$*. When* $I = S$ *we write* P_i *in place of* P_{iS}*.*

For each $i \in \{1,2\}$ and subset I of S recall the notation W_I (introduced in Remark 3.10) of the standard parabolic subgroup of W corresponding to I, it is clear that W_I acts (faithfully) on V_{iI}. This allows us to specify a W_I-action on $\mathrm{Hom}(V_{iI}, \mathbb{R})$ as follows: if $w \in W_I$ and $g \in \mathrm{Hom}(V_{iI}, \mathbb{R})$ then $wg \in \mathrm{Hom}(V_{iI}, \mathbb{R})$ is given by $(wg)v = g(w^{-1}v)$ for all $v \in V_{iI}$. Naturally, when $I = S$ the Coxeter group W acts on $\mathrm{Hom}(V_i, \mathbb{R})$ in a similar way.

Notation 5.2. *For each* $i \in \{1,2\}$ *and subset* I *of* S *we set*

$$P_{iI}^{\#} = \{\, f \in \mathrm{Hom}(V_{iI}, \mathbb{R}) \mid f(x) \geq 0 \text{ for all } x \in P_{iI} \,\},$$

and

$$P_{iI}^{\#\#} = \{\, x \in V_{iI} \mid f(x) \geq 0 \text{ for all } f \in P_{iI}^{\#} \}.$$

Moreover, we set $U_{iI} = \bigcup_{w \in W_I} wP_{iI}^{\#}$*, and write*

$$U_{iI}^{\#} = \{x \in V_{iI} \mid f(x) \geq 0 \text{ for all } f \in U_{iI}\}.$$

When $I = S$ *we write* $P_i^{\#}$*,* $P_i^{\#\#}$*,* U_i *and* $U_i^{\#}$ *in places of* $P_{iS}^{\#}$*,* $P_{iS}^{\#\#}$*,* U_{iS} *and* $U_{iS}^{\#}$ *respectively.*

We call a (convex) subset of a real vector space a *cone* if it is closed under addition and multiplication by positive scalars. It is clear that P_{iI}, $P_{iI}^{\#}$ and $P_{iI}^{\#\#}$ $(i = 1,2)$ are cones for each $I \subseteq S$.

Lemma 5.3. *For each* $i \in \{1,2\}$ *and each* $f \in \mathrm{Hom}(V_i, \mathbb{R})$*, set*

$$\mathrm{Neg}(f) = \{\widehat{x} \in \widehat{\Phi}_i \mid \ x \in \Phi_i^+ \text{ and } f(x) < 0\}.$$

Then $U_i = \{f \in \mathrm{Hom}(V_i, \mathbb{R}) \mid |\mathrm{Neg}(f)| < \infty\}$*.*

Proof. It is enough to prove the U_1 case. Let $f \in U_1$ be arbitrary. Then $f = wg$ for some $w \in W$ and $g \in P_1^{\#}$. Suppose that $x \in \Phi_1^+$ such that $f(x) < 0$. Then $(wg)x = g(w^{-1}x) < 0$. Since $g \in P_1^{\#}$, it follows that $w^{-1}x \in \Phi_1^-$, that is, $\widehat{x} \in N_1(w^{-1})$. Now since $N_1(w^{-1})$ is of finite size, it follows that $U_1 \subseteq \{f \in \mathrm{Hom}(V_1, \mathbb{R}) \mid |\mathrm{Neg}(f)| < \infty\}$. Conversely, suppose that $f \in \mathrm{Hom}(V_1, \mathbb{R})$ with $|\mathrm{Neg}(f)| < \infty$. If $|\mathrm{Neg}(f)| = 0$ then $f \in P_1^{\#} \subseteq U_1$. Thus we may assume that $|\mathrm{Neg}(f)| > 0$, and proceed with an induction. Observe that if $|\mathrm{Neg}(f)| > 0$ then there exists some $s \in S$ such that $f(\alpha_s) < 0$. It is then readily checked that $|\mathrm{Neg}(r_s f)| = |\mathrm{Neg}(f)| - 1$, and hence it follows from our inductive hypothesis that $r_s f \in U_1$. Since U_1 is clearly W-invariant, it follows that $f \in U_1$, and therefore $\{f \in \mathrm{Hom}(V_1, \mathbb{R}) \mid |\mathrm{Neg}(f)| < \infty\} \subseteq U_1$, as required. ☐

The above lemma yields that U_1 and U_2 are cones. In fact, for each $I \subseteq S$ we can show that U_{1I} and U_{2I} are cones. These cones generalize the notion of the Tits cones as defined, for example, in [17, 5.13].

Definition 5.4. We call U_1 and U_2 the *Tits cones* of the Coxeter datum $\mathscr{C}$; and for each $I \subseteq S$ we call U_{1I} and U_{2I} the Tits cones of the the Coxeter datum $\mathscr{C}_I = (\, I, V_{1I}, V_{2I}, \Pi_{1I}, \Pi_{2I}, \langle,\rangle_I \,)$ (where $\langle,\rangle_I$ is the restriction of $\langle,\rangle$ to $V_{1I} \times V_{2I}$).

For each $i \in \{1,2\}$ observe that:

$$\begin{aligned} U_i^{\#} &= \{\, v \in V_i \mid (wf)v \geq 0 \text{ for all } f \in P_i^{\#} \text{ and } w \in W \,\} \\ &= \bigcap_{w \in W} \{\, v \in V_i \mid f(w^{-1}v) \geq 0 \text{ for all } f \in P_i^{\#} \,\} \\ &= \bigcap_{w \in W} \{\, wv \in V_i \mid f(v) \geq 0 \text{ for all } f \in P_i^{\#} \,\} \\ &= \bigcap_{w \in W} wP_i^{\#\#}, \end{aligned} \tag{5.1}$$

and similarly,

$$U_{iI}^{\#} = \bigcap_{w \in W_I} wP_{iI}^{\#\#}, \text{ for each } I \subseteq S. \tag{5.2}$$

The following is a well-known result (a proof can be found in [15, Notes (b) and (c), Lecture 1]).

Lemma 5.5. *Let $I \subseteq S$ be finite. For each $i \in \{1,2\}$, the condition $0 \notin \mathrm{PLC}(\Pi_{iI})$ is equivalent to the existence of an $f_i \in \mathrm{Hom}(V_{iI}, \mathbb{R})$ such that $f_i(x) > 0$ for all $x \in \Pi_{iI}$. Furthermore, $P_{iI}^{\#\#} = P_{iI}$.* □

Lemma 5.6. *For each subset I of S and for each $i \in \{1,2\}$ we have*

$$P_i^{\#\#} \cap V_{iI} \subseteq P_{iI}^{\#\#}.$$

Proof. For each $i \in \{1,2\}$ write $V_i = V_{iI} \oplus V'_{iI}$ where V'_{iI} denotes a (vector space) complement of V_{iI} in V_i. Clearly, every $g_i \in P_{iI}^{\#}$ gives rise to a linear functional g'_i in $P_i^{\#}$ as follows: for any $v \in V_i$ there is a unique decomposition of the form $v = v_{iI} + v'_{iI}$ where $v_{iI} \in V_{iI}$ and $v'_{iI} \in V'_{iI}$, and we simply set $g'_i(v) = g_i(v_{iI})$. Now let $v \in P_i^{\#\#} \cap V_{iI}$, and $f \in P_{iI}^{\#}$ be arbitrary. Then $f(v) = f'(v) \geq 0$ (because $f' \in P_i^{\#}$ and $v \in P_i^{\#\#}$), and hence it follows that $v \in P_{iI}^{\#\#}$, as required. □

Let $x_i \in V_i$ $(i = 1,2)$ be arbitrary. Given condition (C6) of $\mathscr{C}$, it follows that $x_i \in V_{iI}$ for some finite subset I of S. From this fact we may prove the following:

Lemma 5.7. *Suppose that $x_i \in U_i^{\#}$ (for each $i \in \{1,2\}$). Let I be a subset of S such that $x_i \in V_{iI}$. Then $x_i \in U_{iI}^{\#}$. Furthermore, if I is finite then $x_i \in P_{iI}$.*

Proof. Observe that for each $i \in \{1,2\}$,

$$\begin{aligned} x_i &\in U_i^{\#} \cap V_{iI} \\ &\subseteq \bigcap_{w \in W_I} (wP_i^{\#\#} \cap V_{iI}) \\ &= \bigcap_{w \in W_I} w\ (P_i^{\#\#} \cap V_{iI}) && \text{(since } W_I \text{ preserves } V_{iI}\text{)} \\ &\subseteq \bigcap_{w \in W_I} wP_{iI}^{\#\#} = U_{iI}^{\#} && \text{(by Lemma 5.6).} \end{aligned}$$

Finally, it follows from Lemma 5.5 that $x_i \in \bigcap_{w \in W_I} wP_{iI} \subseteq P_{iI}$ whenever I is finite. ☐

Now we are ready for the central result of this section:

Theorem 5.8. *$\langle v_1, v_2 \rangle \leq 0$, for all $v_1 \in U_1^{\#}$ and $v_2 \in U_2^{\#}$.*

Proof. Suppose for a contradiction that there are $v_1 \in U_1^{\#}$ and $v_2 \in U_2^{\#}$ with $\langle v_1, v_2 \rangle > 0$. Replace v_2 by a positive scalar multiple of itself if necessary, we may assume that $\langle v_1, v_2 \rangle = 1$. Let I be a finite subset of S with $v_1 \in V_{1I}$, and let J be a finite subset of S with $v_2 \in V_{2J}$. Next, set $K = I \cup J$. Then Lemma 5.7 yields that $v_1 \in U_{1K}^{\#}$, and $v_2 \in U_{2K}^{\#}$. Since K is finite, it follows from Lemma 5.5 that there are linear functionals $f_1 \in \mathrm{Hom}(V_{1K}, \mathbb{R})$ and $f_2 \in \mathrm{Hom}(V_{2K}, \mathbb{R})$ such that $f_1(\alpha) > 1$ for all $\alpha \in \Pi_{1K}$, and $f_2(\beta) > 1$ for all $\beta \in \Pi_{2K}$. Now set

$$\begin{aligned} \mathscr{A} = \{\, x \in U_{2K}^{\#} \mid f_2(x) \leq f_2(v_2) &\text{ and } \langle z, x \rangle \geq 1 \\ &\text{for some } z \in U_{1K}^{\#} \text{ with } f_1(z) \leq f_1(v_1)\,\}. \end{aligned}$$

Observe that $\mathscr{A} \neq \emptyset$, since $v_2 \in \mathscr{A}$.

Next, put $\epsilon = \frac{2}{|K| f_1(v_1)}$. We claim that for any given $x \in \mathscr{A}$, there exists $y \in \mathscr{A}$ with $f_2(y) \leq f_2(x) - \epsilon$. If we could prove this claim, then starting with $x = v_2$, a finite repetition of this process will produce some $y \in \mathscr{A} \subseteq U_{2K}^{\#} \subseteq P_{2K}$ with $f_2(y)$ being negative, contradicting the fact that $f_2(y) \in f_2(P_{2K}) \subseteq (0, \infty)$. Thus all that remains to do is to prove the above claim. Given arbitrary $x \in \mathscr{A}$, let $z = \sum_{\alpha \in \Pi_{1K}} \lambda_\alpha \alpha \in U_{1K}^{\#}$ be such that $\langle z, x \rangle \geq 1$, $f_1(z) \leq f_1(v_1)$, and $\lambda_\alpha \geq 0$ for all $\alpha \in \Pi_{1K}$. Note that these conditions imply that $\sum_{\alpha \in \Pi_{1K}} \lambda_\alpha \langle \alpha, x \rangle \geq 1$, which in turn implies that $\lambda_{\alpha_{s_0}} \langle \alpha_{s_0}, x \rangle \geq \frac{1}{|K|}$ for some $s_0 \in K$. Then

$$\langle \alpha_{s_0}, x \rangle \geq \frac{1}{\lambda_{\alpha_{s_0}} |K|} \geq \frac{1}{f_1(v_1)|K|} = \frac{\epsilon}{2},$$

since $\lambda_{\alpha_{s_0}} \leq f_1(z) \leq f_1(v_1)$. Set $y = r_{s_0} x$. Observe that (5.2) indicates that $U_{2K}^{\#}$ is W_K-invariant, and given that $r_{s_0} \in W_K$, we have $y \in U_{2K}^{\#}$. Moreover,

$$f_2(y) = f_2(x) - 2\langle \alpha_{s_0}, x \rangle f_2(\beta_{s_0}) < f_2(x) - \epsilon < f_2(v_2).$$

Thus to establish our claim, we only need to show that $y \in \mathscr{A}$, and this in turn amounts to finding some $t \in U^{\#}_{1K}$ such that $\langle t, y\rangle \geq 1$, and $f_1(t) \leq f_1(v_1)$. First, suppose that $\langle z, \beta_{s_0}\rangle \geq 0$. Put $t = r_{s_0}z$. Since $U^{\#}_{1K}$ is W_K-invariant and $z \in U^{\#}_{1K}$, it follows that $t \in U^{\#}_{1K}$. Moreover,

$$\langle t, y\rangle = \langle r_{s_0}z, r_{s_0}x\rangle = \langle z, x\rangle \geq 1$$

and

$$f_1(t) = f_1(z) - 2\langle z, \beta_{s_0}\rangle f_1(\alpha_{s_0}) \leq f_1(z) \leq f_1(v_1),$$

thus proving $y \in \mathscr{A}$ in the case $\langle z, \beta_{s_0}\rangle \geq 0$. Next, suppose that $\langle z, \beta_{s_0}\rangle < 0$. Then $t = z$ will do, indeed,

$$\begin{aligned} \langle z, y\rangle &= \langle z, x - 2\langle \alpha_{s_0}, x\rangle \beta_{s_0}\rangle \\ &= \langle z, x\rangle - 2\langle \alpha_{s_0}, x\rangle\langle z, \beta_{s_0}\rangle \\ &\geq \langle z, x\rangle \\ &\geq 1, \end{aligned}$$

and by our construction, $z \in U^{\#}_{1K}$ and $f_1(z) \leq f_1(v_1)$, thus proving $y \in \mathscr{A}$ in the case $\langle z, \beta_{s_0}\rangle < 0$ too. This completes the proof of the claim, and hence the theorem follows. ☐

6. Acknowledgments

A few results presented in this paper are taken from the author's PhD thesis [10] and the author wishes to thank A/Prof. R. B. Howlett for all his help and encouragement throughout the author's PhD candidature. The author also wishes to thank Prof. G. I. Lehrer and Prof. R. Zhang for supporting this work.

References

1. A. Björner and F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics, GTM 231, Springer, 2005
2. N. Bourbaki, *Groupes et algebras de Lie, Chapitres 4, 5 et 6* , Hermann, Paris, 1968
3. B. Brink and R. B. Howlett, *A finiteness property and an automatic structure of Coxeter groups*, Math. Ann. **296** (1993), 179–190.
4. B. Brink, *The set of dominance-minimal roots*, J. Algebra, **206** (1998), 371–412.
5. F. Bergeron, N. Bergeron, R.B. Howlett and D.E. Taylor *A Decomposition of the Descent Algebra of a Finite Coxeter Group*, Journal of Algebraic Combinatorics, **1** (1992), 23–44
6. P. E. Caprace and B. Rémy, *Groups with a root group datum*, Innov. Incidence Geom. 9 (2009), 5–77.
7. V. Deodhar, *On the root system of a Coxeter group*, Comm. Algebra, 10(6): 611–630, 1982.
8. M. Dyer, *Hecke algebras and reflections in Coxeter groups*, PhD thesis, University of Sydney, 1987.
9. M. Dyer, *Reflection Subgroups of Coxeter Systems*, J. Algebra, **135** (1990), 57–73.
10. X. Fu, *Root systems and reflection representations of Coxeter groups*, PhD thesis, University of Sydney, 2010.

11. X. Fu, *The dominance hierarchy in root systems of Coxeter groups*, J. Algebra, **366** (2012), 187–204.
12. J.-Y. Hée, *Système de racines sur un anneau commutatif totalement ordonné*, Geom. Dedicata, **37**, (1991), 65–102.
13. H. Hiller, *Geometry of Coxeter Groups*. Research Notes in Mathematics 54, Pitman (Advanced Publishing Program), Boston-London, 1981.
14. R. B. Howlett, *Normalizers of parabolic subgroups of reflection groups*, J. London Math. Soc. (2) 21 (1980), no. 1, 62–80.
15. R. B. Howlett, *Introduction to Coxeter groups*, Lectures given at ANU, 1996 (available at http://www.maths.usyd.edu.au/res/Algebra/How/1997-6.html).
16. R. B. Howlett, P. J. Rowley and D. E. Taylor, *On Outer Automorphism groups of Coxeter Groups*, Manuscripta Math. 93 (1997), 499–513.
17. J. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Stud. Adv. Math. vol. 29, Cambridge Univ. Press, 1990.
18. V. G. Kac, *Infinite-dimensional Lie algebras*, third edition, Cambridge University Press, Cambridge, 1990.
19. D. Krammer, *The conjugacy problem for Coxeter groups*, PhD thesis, Universiteit Utrecht, 1994.
20. G. A. Maxwell, *Sphere packings and hyperbolic reflection groups*, Journal of Algebra, **79**, (1982), 78–97.
21. R. V. Moody and A. Pianzola, *On infinite root systems*, Trans. Amer. Math. Soc. **315** (1989), 661–696.
22. È. B. Vinberg, *Discrete Linear Groups Generated by Reflections*, Izv. Akad. Nauk SSSR Ser. Mat. **35** (1971), 1072–1112.